



Contents lists available at SciVerse ScienceDirect

Biochimica et Biophysica Acta

journal homepage: www.elsevier.com/locate/bbabbio

Review

Photosystem II and the unique role of bicarbonate: A historical perspective [☆]

Dmitriy Shevela ^{a,*}, Julian J. Eaton-Rye ^b, Jian-Ren Shen ^c, Govindjee ^{d,e,f,g,**}

^a Centre for Organelle Research, University of Stavanger, Kristine Bonnevis vei 22, N-4036 Stavanger, Norway

^b Department of Biochemistry, University of Otago, PO Box 56, Dunedin 9054, New Zealand

^c Division of Bioscience, Graduate School of Natural Science and Technology/Faculty of Science, Okayama University, Okayama 700-8530, Japan

^d Department of Plant Biology, University of Illinois at Urbana-Champaign, 265 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801, USA

^e School of Life Sciences, Jawaharlal Nehru University, New Delhi, 110067, India

^f Department of Biochemistry, University of Illinois at Urbana-Champaign, 265 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801, USA

^g Center of Biophysics & Computational Biology, University of Illinois at Urbana-Champaign, 265 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801, USA

ARTICLE INFO

Article history:

Received 14 January 2012

Received in revised form 5 April 2012

Accepted 6 April 2012

Available online xxx

Keywords:

Bicarbonate (hydrogen carbonate) effect

Electron transport

Protonation reaction

O₂ evolution

Photosystem II

ABSTRACT

In photosynthesis, cyanobacteria, algae and plants fix carbon dioxide (CO₂) into carbohydrates; this is necessary to support life on Earth. Over 50 years ago, Otto Heinrich Warburg discovered a unique stimulatory role of CO₂ in the Hill reaction (i.e., O₂ evolution accompanied by reduction of an artificial electron acceptor), which, obviously, does not include any carbon fixation pathway; Warburg used this discovery to support his idea that O₂ in photosynthesis originates in CO₂. During the 1960s, a large number of researchers attempted to decipher this unique phenomenon, with limited success. In the 1970s, Alan Stemler, in Govindjee's lab, perfected methods to get highly reproducible results, and observed, among other things, that the turnover of Photosystem II (PSII) was stimulated by bicarbonate ions (hydrogen carbonate): the effect would be on the donor or the acceptor, or both sides of PSII. In 1975, Thomas Wydrzynski, also in Govindjee's lab, discovered that there was a definite bicarbonate effect on the electron acceptor (the plastoquinone) side of PSII. The most recent 1.9 Å crystal structure of PSII, unequivocally shows HCO₃⁻ bound to the non-heme iron that sits in-between the bound primary quinone electron acceptor, Q_A, and the secondary quinone electron acceptor Q_B. In this review, we focus on the historical development of our understanding of this unique bicarbonate effect on the electron acceptor side of PSII, and its mechanism as obtained by biochemical, biophysical and molecular biological approaches in many laboratories around the World. We suggest an atomic level model in which HCO₃⁻/CO₃²⁻ plays a key role in the protonation of the reduced Q_B. In addition, we make comments on the role of bicarbonate on the donor side of PSII, as has been extensively studied in the labs of Alan Stemler (United States) and Vyacheslav Klimov (Russia). We end this review by discussing the uniqueness of bicarbonate's role in oxygenic photosynthesis and its role in the evolutionary development of O₂-evolving PSII. This article is part of a Special Issue entitled: Photosynthesis Research for Sustainability: from Natural to Artificial Photosynthesis.

© 2012 Published by Elsevier B.V.

1. Introduction

1.1. Role of inorganic carbon and its interconversion in living organisms

Carbon dioxide (CO₂) is not only a greenhouse gas in the Earth's atmosphere, but also a key metabolite in living organisms, where it plays an essential role in such fundamental biological processes as respiration and

photosynthesis. Due to its ability to exist in equilibrium with carbonic acid (H₂CO₃) and bicarbonate (HCO₃⁻, IUPAC's recommended term is *hydrogen carbonate*, but in this review we use its traditional and well-known term '*bicarbonate*') (see Fig. 1); CO₂ produced by cells during aerobic metabolism of glucose and fats provides the acid (H⁺ and CO₂) and base (HCO₃⁻) components for the so-called *bicarbonate buffering system*. This buffering system maintains both intracellular and extracellular pH.

Abbreviations: CA, carbonic anhydrase; Chl, chlorophyll; DCMU, 3-(3,4-dichlorophenyl)-1,1-dimethylurea (diuron); DPC, 1,5-diphenylcarbazine; HCO₃⁻, hydrogen carbonate (bicarbonate) ion; MS, mass spectrometry; NHI, non-heme iron; OEC, oxygen-evolving complex; P680, primary electron donor molecule (Chl) in Photosystem II; P700, primary electron donor molecule (Chl) in Photosystem I; Pheo, pheophytin; PQ, plastoquinone; PQH₂, plastoquinol; PSI, Photosystem I; PSII, Photosystem II; Q_A, primary quinone electron acceptor of PSII; Q_B, secondary quinone electron acceptor of PSII; RC, reaction center; S_i, redox state of the OEC, where *i* is the number of stored oxidizing equivalents; TL, thermoluminescence

[☆] This article is part of a Special Issue entitled: Photosynthesis Research for Sustainability: from Natural to Artificial Photosynthesis.

* Corresponding author at: Centre for Organelle Research, University of Stavanger, Kristine Bonnevis vei 22, N-4036 Stavanger, Norway. Tel.: +47 518 31810; fax: +47 518 31860.

** Corresponding author at: Department of Plant Biology, University of Illinois at Urbana-Champaign, 265 Morrill Hall, 505 South Goodwin Avenue, Urbana, IL 61801, USA. Tel.: +1 217 337 0627; fax: +1 217 244 7246.

E-mail addresses: dmitry.shevela@uis.no (D. Shevela), gov@illinois.edu (Govindjee).

0005-2728/\$ – see front matter © 2012 Published by Elsevier B.V.

doi:10.1016/j.bbabbio.2012.04.003

Please cite this article as: D. Shevela, et al., Photosystem II and the unique role of bicarbonate: A historical perspective, *Biochim. Biophys. Acta* (2012), doi:10.1016/j.bbabbio.2012.04.003

The interconversion of inorganic carbon, on the other hand, allows rapid transport of its species ($\text{CO}_2/\text{HCO}_3^-/\text{CO}_3^{2-}$) in all cells. While HCO_3^- is poorly soluble in biological membranes, CO_2 can freely diffuse in and out of the cell. Therefore, $\text{HCO}_3^- \rightarrow (\text{H}_2\text{CO}_3) \rightarrow \text{CO}_2$ interconversion facilitates the transport of inorganic carbon in the form of CO_2 into intracellular space, while the reversed conversion ($\text{CO}_2 \rightarrow (\text{H}_2\text{CO}_3) \rightarrow \text{HCO}_3^-$) provides trapping of the CO_2 within the cell in the form of HCO_3^- . Although the reversible hydration of CO_2 and dehydration of HCO_3^- occurs spontaneously, even in the absence of catalysts, most—if not all—organisms have Zn-containing *carbonic anhydrases* (CAs) that catalyze this ubiquitous conversion. By speeding up these reactions (k_{cat} can reach $\sim 1 \times 10^6 \text{ s}^{-1}$), CAs play an essential role in a wide range of biochemical and physiological processes [1].

All photosynthetic organisms need atmospheric CO_2 to store harvested energy from sunlight in the form of energy-rich carbohydrates. However, in the oxygenic photosynthesizers (cyanobacteria, algae and higher plants), CO_2 is not only required as the terminal electron acceptor to synthesize carbohydrates, but also for the regulation of photosynthetic electron transport in Photosystem II (PSII), the enzyme responsible for light-induced primary charge separation and subsequent water oxidation [2]. The latter is known as the ‘bicarbonate effect’. This review summarizes historical discoveries related to the ‘bicarbonate effect’ and outlines our current state of knowledge about the location and role of HCO_3^- in PSII.

1.2. The ‘bicarbonate effect’ and Otto Heinrich Warburg

Despite the fact that the reduction of CO_2 to carbohydrates within the Calvin–Benson cycle is driven by the products of the light reactions of photosynthesis, ATP and NADPH, it does not directly require light, and thus, belongs to the photosynthetic light-independent (‘dark’) reactions [3]. Until the discovery (1958) of the ‘bicarbonate effect’ on the light-dependent electron flow by Otto Warburg and Günter Krippahl, CO_2 was assumed to be involved only in the ‘dark’ and not in the ‘light’ reactions [4]. Warburg and Krippahl found that the high rates of the Hill reaction (the reaction which allows the study of electron flow in isolated broken chloroplasts illuminated in the presence of an artificial electron acceptor *via* the measurements of O_2 production) required the presence of CO_2 in the gas phase above the sample suspension. Specifically, it was shown that the O_2 evolution rate measured in grana isolated from kohlrabi leaves in the presence of quinone (as electron acceptor) was significantly higher when argon atmosphere above the sample contained 1.4% CO_2 (v/v) (see Fig. 2). In spite of the earlier difficulties with its reproducibility, many research groups confirmed this phenomenon (outlined in Section 3.1). Later on, Alan Stemler and Govindjee [5] significantly improved reproducibility of the bicarbonate effect by developing a reliable method of $\text{CO}_2/\text{HCO}_3^-$ depletion.

Otto Warburg believed that the observed phenomenon provides evidence for his ‘photolyte theory’, in which O_2 originates from the splitting of ‘activated CO_2 ’, not from water. In 1964, he noted “As was expected, no proof of water photolysis survived the discovery of ‘active CO_2 ’” [6]. Despite this mistaken interpretation, the finding made by Warburg and Krippahl was fundamental to subsequent research of the ‘bicarbonate effect’ on light-induced electron transport

during photosynthesis. Their discovery initiated long-term debates about possible action site(s) and role(s) of inorganic carbon on photosynthetic O_2 production. Thus, intensive studies by many laboratories explored the possibility that HCO_3^- (CO_3^{2-}) (and not CO_2) is required for both PSII electron transport efficiency and for the photo-assembly of the inorganic core (the Mn_4CaO_5 cluster) of the O_2 -evolving complex (OEC) of PSII (see Sections 2 and 3; for previous historical overviews, see [7–14]).

1.3. Photosystem II and the sites of the ‘bicarbonate effect’

PSII is a large multi-component pigment-protein complex, which is incorporated into the thylakoid membrane of all oxygenic photosynthetic organisms (for reviews on PSII, see [2,15]). Fig. 3 shows a schematic view of PSII in higher plants and green algae and its important redox cofactors, which are thought to be the same as in cyanobacteria (for further details on the cyanobacterial PSII structure, see [16] and [17]). PSII acts as a water:plastoquinone oxidoreductase, catalyzing the following reaction:



Thus, the light-induced charge separation between the reaction center (RC) chlorophyll (Chl) molecules in the D1 protein (Chl_{D1} and P680) and pheophytin (Pheo_{D1}), and the formation of the stabilized radical ion pair $\text{P680}^{+\bullet}\text{Pheo}^{-\bullet}$ (for reviews, see [18,19]), lead to two reactions: (1) water splitting (oxidation) to O_2 , protons and electrons on the luminal side of PSII, the so-called *electron donor side* of PSII with $\text{P680}^{+\bullet}$ as the driving force and, (2) the reduction of plastoquinone (PQ) to plastoquinol (PQH_2) on the stromal, the *electron acceptor side*, of PSII with proton uptake and $\text{QA}^{\bullet-}$ acting as the reductant.

Extensive data show that HCO_3^- ions, under appropriate experimental conditions, have effects on both the acceptor side and the donor side reactions of PSII (Fig. 3). The focus of this review is the bicarbonate effects related to the electron acceptor side, where HCO_3^- is known to bind (see Fig. 3 and Section 3.2 for the current model) and to play an important role in facilitating the reduction of Q_B , and in protonation reactions near the Q_B -site. We, however, briefly discuss here, for completeness, the possible roles of HCO_3^- on the ‘donor side’ reactions of PSII. For historical surveys on the discoveries of the HCO_3^- effect on the PSII donor side, see several references [9,10,14,20–22].

2. Bicarbonate and the donor side of Photosystem II

The role of HCO_3^- on the donor side reactions of PSII has been extensively studied by many researchers, but mainly in the laboratories of Alan Stemler (University of California, USA) and Vyacheslav Klimov (Institute of Basic Biological Problems, Russia), as mentioned earlier.

In the early 1970s, Stemler with co-workers [5,23] were the first to propose the water-oxidizing side of PSII as a possible site for the HCO_3^- effect (Fig. 4). However, in 1975, Thomas Wydrzynski and Govindjee obtained evidence for the participation of HCO_3^- ions in the electron transfer kinetics on the acceptor side of PSII [24] (Fig. 5). This discovery was supported by numerous subsequent experiments (see




Fig. 1. Conversion of inorganic carbon species including acid–base ionization/dissociation constant (pK_a) values for hydrogen carbonate (bicarbonate) anion. See text for further details.




Fig. 2. CO₂ (HCO₃⁻) effect on the rate of the Hill reaction as was found by Otto Warburg and Günter Krippahl [4] in isolated kohlrabi grana suspended in 0.1% KCl. The measurements were performed in the presence of 2.1 mg quinone as electron acceptor under argon (closed symbols) or argon + 1.4% CO₂ (v/v) (open symbols) in the gas phase.

165 Section 3), and later on, the non-heme iron (NHI) between Q_A and Q_B
 166 was shown to play an essential role in HCO₃⁻ binding [12,25]. On the
 167 other hand, some (among them Helmut Metzner, Werner Kreutz, and
 168 Alan Stemler) believed that HCO₃⁻ may act as a substrate or a chemical
 169 intermediate in photosynthetic O₂ evolution, possibly coupled with CA
 170 activity [26–28]. Thus, Stemler and collaborators continued to investigate
 171 the possible involvement of HCO₃⁻ ions in the mechanism of O₂ evolution
 172 on the oxidizing side of PSII (reviewed in [9,22]). Stemler's reports, as well
 173 as reports of some others [29,30] indicated that HCO₃⁻ may affect both the
 174 electron acceptor and donor sides of PSII. Undoubtedly, however, the dis-

covery of the 'acceptor-side' effect inadvertently affected the search for
 175 specific effects of HCO₃⁻ on the donor side, and inevitably led to a contro-
 176 versy on the interpretation of the 'bicarbonate effects'.
 177

178 Since the mid 1990s, the idea for an additional role of HCO₃⁻ on the
 179 electron donor side of PSII was revived by a series of experiments per-
 180 formed in the laboratory of Vyacheslav Klimov. The studies by Klimov
 181 and collaborators indicated that HCO₃⁻ ions are required for (1) the
 182 efficient photo-induced assembly of the Mn₄CaO₅ cluster capable of
 183 water splitting, (2) the stability of the OEC, and (3) the protection
 184 of the donor side of PSII against photoinhibition and thermoinactiva-
 185 tion (reviewed in [10,14]). Other groups (see, for instance, [31–35])
 186 also obtained indication for the requirement of HCO₃⁻ on the water-
 187 splitting side of PSII. However, the binding site(s) and the role(s) of
 188 HCO₃⁻ ions in the water-splitting reaction of PSII remain unclear
 189 (and, therefore, appear 'questionable' to the authors; see Fig. 3). The
 190 following main proposals for the involvement of HCO₃⁻ in the events
 191 on the water-oxidizing side of PSII have been considered:

- (i) Exchangeable HCO₃⁻ is an intermediate substrate for photosynthetic
 192 water oxidation; water is delivered to the Mn₄CaO₅ cluster in the
 193 form of HCO₃⁻ (or peroxidicarbonic acid; H₂C₂O₆). Initially proposed
 194 by Helmut Metzner [27] as an alternative to Warburg's 'photolyte
 195 theory' (mentioned above) and later elaborated by Alan Stemler,
 196 [26] and by Paul Castelfranco with co-authors [36], this hypothe-
 197 sis has become obsolete, in our opinion, because of various studies
 198 using isotope ratio mass spectrometry (MS) in combination with
 199 ¹⁸O-labeling of H₂O and HCO₃⁻ [37–41], UV spectrophotometry
 200 under high backpressure of CO₂ [38], and light-induced FT-IR dif-
 201 ference spectroscopy [42].
 202




Fig. 3. Schematic representation of PSII in higher plants and green algae (only core proteins are shown) and two sites (acceptor and donor) where bicarbonate (HCO₃⁻; hydrogen carbonate) has effects. While the acceptor side bicarbonate is known to bind to the NHI (Fe²⁺) between Q_A and Q_B, the exact location of the donor side bicarbonate is unknown. The acceptor side bicarbonate may also be bound to the NHI in the form of carbonate (CO₃²⁻). In cyanobacteria the sites of HCO₃⁻ effects are the same, but some components of PSII are different (for further details see [15]). The pathway of the electron flow through PSII is shown by black arrows. Other abbreviations: D1 and D2, the reaction center proteins; P680, the reaction center Chl molecule; Chl_{D1}, the primary electron donor on D1; Pheo_{D1}, the primary electron acceptor on D1 (pheophytin); Chl_{D2} and Pheo_{D2}, symmetrically related cofactors on D2 (inactive branch; do not participate in linear electron transfer through PSII); Mn₄CaO₅, inorganic core of the OEC; Y_Z (on D1) and Y_D (on D2), the redox active tyrosine residues; PQ, mobile plastoquinone molecule; CP43 and CP47, Chl-protein complexes of 43 and 47 kDa; LHC-II, light-harvesting complex II; PsaO (33 kDa), PsaP (23 kDa) and PsaQ (17 kDa), extrinsic proteins of PSII; Cyt b559, redox active cytochrome b559.




Fig. 4. O₂ yield obtained on the third flash (Y_3) as a function of the dark-time between the second and the third flash (Δt_{23}) as measured by Stemler et al. [23] in dark-adapted HCO₃⁻-depleted chloroplast suspensions in the presence and the absence of bicarbonate. The frequency of the main flash train was 1 Hz. Y_3 values were normalized with respect to the steady-state O₂ yield. Open triangles: HCO₃⁻-depleted chloroplast suspensions were injected onto the Pt electrode to final Chl concentration of 0.3 mg ml⁻¹. The measurements were performed in buffered medium, which contained 0.25 M NaCl, 0.04 M Na acetate, 0.05 M Na phosphate buffer (pH 6.8), 20 μg ml⁻¹ of ferredoxin, and 0.5 mM NADP⁺. Closed circles: the same as above but after re-addition of 10 mM NaHCO₃. Adapted and modified from [23].

(ii) *Non-exchangeable, tightly bound HCO₃⁻ being a structural part of the Mn₄CaO₅ cluster may alter the redox properties of the Mn cations, and thus, is required for the functionality and stability of the assembled OEC.* The studies carried out by Klimov and collaborators showed stabilizing and protective effects of HCO₃⁻ on the donor (water-oxidizing) side of PSII [43–50]. One of the interpretations of the




Fig. 5. First experimental evidence for the action of bicarbonate on the electron acceptor side of PSII reported by Wydrzynski and Govindjee in 1975 [24]. A comparison of variable Chl *a* fluorescence on concentration of HCO₃⁻ and an inhibitor of PSII DCMU. (A) HCO₃⁻-depleted chloroplasts at various concentrations of NaHCO₃. (B) Non-HCO₃⁻-depleted chloroplasts at various concentrations of DCMU. Before the measurements, the samples were incubated in the dark for 5 min. Note that the fluorescence induction curves obtained under HCO₃⁻-free conditions look like the one obtained after addition of 10⁻⁶ M DCMU. Fluorescence was measured at 685 nm upon excitation with a broad blue light at a Chl concentration of 12.5 μg ml⁻¹. Modified and adapted from [24].

observed effects was the idea that HCO₃⁻ may function as a ligand to the Mn₄CaO₅ cluster or an integral cofactor of the OEC [10]. In addition, in the PSII crystal structure by Ferreira et al. [51] at a resolution of 3.5 Å, HCO₃⁻ (or CO₃²⁻) anion was tentatively included as a ligand bridging Mn and Ca ions within the OEC. However, the latter could not be supported by the most recent X-ray crystallography studies of PSII at higher resolutions [16,17,52]. At the same time, all these crystallographic studies clearly displayed HCO₃⁻ as a ligand of the NHI between Q_A and Q_B (for further details, see Section 3.2). Earlier sensitive differential infrared gas analyzer and MS measurements [7,53] also clearly showed only ~1 HCO₃⁻/CO₂ molecule bound per PSII RC (see Section 3.1.2.3). Further, a recent re-examination of the structural coupling of HCO₃⁻ to the OEC by FT-IR spectroscopy provided no indication for any HCO₃⁻-bands from the OEC during the S-state transitions [42]. This is also consistent with the results obtained by flash-induced O₂ evolution pattern (FIOP) studies, where the redox potentials of the S states of the OEC were found to be unaffected by HCO₃⁻ depletion *via* washing with CO₂/HCO₃⁻-free buffer [54]. Moreover, evidence for the absence of tightly bound HCO₃⁻ in the first coordination sphere of the Mn₄CaO₅ cluster was obtained by isotope ratio MS [41] (for details, see Section 3.1.3.3) and GC-MS [55] studies. In addition, HCO₃⁻, as a structural part of the OEC has not been supported by the computational models based on density functional theory (DFT) and quantum mechanics/molecular mechanics (QM/MM) studies [56,57]. It is, therefore, very unlikely that HCO₃⁻ is a ligand or strongly coupled cofactor to the Mn₄CaO₅ cluster in its assembled state.

(iii) *Acting as a transient ligand to Mn ions, HCO₃⁻ is a native cofactor in the photo-assembly (photo-activation) process of the Mn₄CaO₅ cluster that assembles in the OEC-depleted PSII centers that are free of inorganic cofactors, but HCO₃⁻ is not part of the assembled cluster.* This suggestion is based on the results obtained by Klimov and co-workers [10] demonstrating a pronounced stimulating effect of HCO₃⁻ ions on the electron donation from exogenous Mn²⁺ ions to Mn-depleted PSII and the photo-induced reconstitution of the functional OEC [43–45,48,58]. Further experiments in collaboration with the group of Charles Dismukes provided evidence for the requirement of HCO₃⁻ (CO₃²⁻) for *in vitro* light-driven assembly of the Mn₄CaO₅ cluster (for details, see [31,59]; for reviews on the photo-assembly of the OEC, see [60,61]). Although electrochemical characterizations of Mn-HCO₃⁻ complexes [62–64] as well as electron paramagnetic resonance/electron spin echo envelope modulation (EPR/ESEEM) spectroscopy studies of assembly intermediates [65,66] strongly support this idea, there is no experimental data demonstrating a HCO₃⁻ requirement for the assembly process *in vivo*.

(iv) *HCO₃⁻ indirectly stabilizes the OEC by binding to extrinsic proteins or some other protein components of PSII in the vicinity of the Mn₄CaO₅ cluster.* Pobeguts et al. [67] demonstrated a protective effect of HCO₃⁻ against extraction of the extrinsic proteins (especially PsbO—the Mn-stabilizing protein) of the OEC after treatment of pea PSII membrane fragments with urea. Moreover, the specific high-affinity binding of HCO₃⁻ (or CO₂) to the PsbO protein has been proposed based on the recent observations of HCO₃⁻-dependent re-arrangements in the PsbO protein [68]. However, as mentioned above, no HCO₃⁻ was detected bound on the protein components belonging to the donor side of PSII of thermophilic cyanobacteria by Umena et al. [17] in their recent crystal structure at 1.9 Å resolution. Nevertheless, since significant differences are known to exist between proteins of cyanobacteria and plants (reviewed in [69,70]), the possibility of HCO₃⁻ binding to protein components in higher plants needs to be addressed by future experiments.

(v) *Mobile, exchangeable HCO₃⁻ is involved in proton removal during photosynthetic water oxidation; it may work coupled with the*

PSII-donor-side-associated CA. Deprotonation reactions and removal of protons away from the OEC are thought to have significant impact on the thermodynamics of water splitting [71]. Ananyev et al. [34] proposed that HCO_3^- may play an indirect role in water splitting as a proton transfer mediator and recent results support this proposal [32,35]. In fact, such an interpretation of the bicarbonate effect on the water-oxidizing side of PSII may explain a large body of existing data, especially in connection with the PSII-associated CA activity, which was experimentally shown in many studies [72–78]. Stemler, based on circumstantial evidence, was the first to suggest that a thylakoid CA might be involved in the ‘donor-side’ effects of HCO_3^- [74,79–81]. Experiments by Shutova et al. [32] show that in *Chlamydomonas* (*C.*) *reinhardtii*, both HCO_3^- and Cah3 (the CA protein in *C. reinhardtii* associated with the PSII donor side) have specific ‘donor-side’ effects on proton release steps, but not on electron transfer. Moreover, there are also some indications for a similar role of CA and HCO_3^- in higher plants, although both CA and HCO_3^- requirements were found to be lower than that observed in *C. reinhardtii* [82]. Shutova et al. [32] suggested that a CA/ HCO_3^- system in *C. reinhardtii* may facilitate proton removal away from the OEC during water splitting by accelerating interconversion between HCO_3^- and CO_2 (see Fig. 1). Indeed, if the luminal “working” pH under illumination is 5.4–5.7, as shown recently [83,84], one can assume that due to the strong deficit of HCO_3^- species at this pH range, the presence of CA activity is ‘naturally’ required for the fast production of these species from CO_2 .

We note here, that to our knowledge, most of the above bicarbonate-related investigations of cyanobacteria, algae, and higher plants, except for a few with intact alga *Chlamydomonas stellata* [29] and the hypercarbonate-requiring cyanobacterium *Arthrospira maxima* [85,86] have been limited to *in vitro* studies of isolated thylakoids, PSII membrane fragments and PSII particles. Therefore, we emphasize that the effect and the function of HCO_3^- on the donor side of assembled PSII may be different (if any) when the protein environment is intact as to when it is disrupted, e.g., as a result of sample preparation. Thus, further research is needed to study this option and to elucidate the role of HCO_3^- on the water-oxidizing side of PSII.

3. Bicarbonate and the acceptor side of Photosystem II

In contrast to what little is known regarding the effect of bicarbonate on the donor side of PSII, we know a great deal about the role of HCO_3^- on the acceptor-side of PSII—and its binding is obvious in the high resolution structures of cyanobacterial PSII RCs. Therefore, the remaining historical perspective in this review will focus on the research efforts related to the discoveries of the bicarbonate effect on the electron flow within PSII, and the HCO_3^- -dependent regulation of electron transport on the acceptor side. Hence, the current state of our knowledge about the location and the function of HCO_3^- is also discussed in detail.

3.1. Time-line of discoveries on HCO_3^- in the electron flow of Photosystem II

3.1.1. The early work: from Otto Warburg to Norman Good

In 1948, Boyle [87] had observed that O_2 evolution by ground-up spinach leaves, when *p*-benzoquinone was added, was absent when KOH was included (to absorb CO_2) in the center well of a manometer vessel; thus, Boyle concluded that CO_2 was necessary for the benzoquinone Hill reaction. Although the conclusion was confirmed by Warburg and Krippahl in 1960 [88], Boyle’s results were artifacts as suggested by Warburg and Krippahl (1958) [4] and, as shown, in 1961, by Abeles et al. [89]: benzoquinone in the main

vessel distilled into the KOH-soaked filter paper in the center well and the mixture consumed O_2 balancing O_2 evolution from the broken leaves. The discovery of the ‘bicarbonate effect’ by Warburg and Krippahl has already been mentioned in Section 1.2.

- In 1961, Abeles et al. [89] confirmed Warburg and Krippahl’s results in kohlrabi chloroplasts, i.e., requirement of CO_2 for the Hill reaction. Warburg’s idea that O_2 arose from CO_2 (see Section 1.2) had to be tested. For this purpose, Abeles et al. used MS that distinguishes O_2 evolution and metabolism of CO_2 . They observed changes only in O_2 release and none in CO_2 metabolism. Thus, Warburg’s idea was not supported. However, Abeles and co-workers could not reproduce this effect in sugar beet chloroplasts leading them to conclude that the effect was not universal.
- During 1960–1962, Stern and Vennesland [90,91] observed that the ferricyanide-supported Hill reaction, in spinach and kohlrabi chloroplasts suspended in buffered media, declined much faster, with time, with CO_2 present. Addition of CO_2 restored Hill activity. Further, in 1963, Vennesland, who was still supporting Warburg’s point of view, reported stimulation of the Hill reaction with different electron acceptors, using thylakoids from various plant sources [92].
- In 1962, Izawa [93] introduced the use of CA to the reaction medium, while CO_2 was being removed; this hastened the time of CO_2 -depletion and gave much more reliable results; Izawa found larger effects in broken than in intact chloroplasts.
- In 1963, Heise and Gaffron [94] reported decreases in O_2 evolution during the Hill reaction with *p*-benzoquinone in the cyanobacterium *Anacystis nidulans* (*Synechococcus elongatus* strain PCC 7942) and in the green alga *Scenedesmus obliquus* (strain D3) in the absence of CO_2 . However, these authors suggested that this effect is not an important one since many different metabolic reactions have been shown to be dependent on traces of CO_2 .
- During 1963–1965, Good [95,96] discovered that CO_2 dependence of the Hill reaction, in pea chloroplasts, was highly influenced by the addition of anions, particularly of formate and acetate; none of the anions used could act as bicarbonate; thus, bicarbonate was considered to have a specific stimulatory effect in electron transport during the Hill reaction; uncouplers of phosphorylation had no effect on electron transport in CO_2 -depleted chloroplasts.
- During 1964 and 1965, Punnett and Iyer [97], Punnett [98] and Batra and Jagendorf [99] discovered that in addition to the effects of $\text{CO}_2/\text{HCO}_3^-$ on electron transport, an additional, although a different effect, exists on photophosphorylation. In their 1978 review on the bicarbonate effect, Govindjee and Van Rensen [11] have called this separate effect, the “Punnett Effect”; however, it will not be discussed further in this review since CO_2 was not a requirement for phosphorylation, whereas it is a requirement for electron transport.
- In 1967, West and Hill [100] confirmed the existence of the stimulatory role of CO_2 in both dichlorophenol indophenol (DCPIP) and ferricyanide Hill reactions in pea chloroplasts, and as Izawa had stated, the effect was larger in broken, than in intact, chloroplasts.

3.1.2. Work at the University of Illinois at Urbana-Champaign (UIUC): from Alan Stemler to Jin Xiong and collaborations with other groups

Most of the research up to this point was aimed to see if there was an effect of CO_2 on the Hill reaction, i.e., electron flow from water to NADP^+ (see a review [11]). There were, in general, considerable variations in the magnitude of the effects; further, the conditions producing them showed considerable differences. Govindjee, one of the authors of this review, presented a lecture to a graduate level course in late 1960s or early 1970s, where he talked about this effect emphasizing the *out-of-this-world* ideas of Otto Warburg—that this effect implies that O_2 comes from CO_2 . To the surprise of Govindjee, one of his own doctoral students in the class wanted to pursue this as his PhD thesis project. Govindjee attempted to discourage such an undertaking as it was very risky, but then the student Alan Stemler persisted. The rest is history. We present below a time line of research from 1973 to 1998 in Govindjee’s laboratory in this area of research (also see a different perspective in Stemler’s reviews

[9,21]). We define the ‘bicarbonate effect’ as follows: addition of bicarbonate to $\text{CO}_2/\text{HCO}_3^-$ -depleted samples restores fully the electron transport to that without the depletion process.

3.1.2.1. 1970s: research at the UIUC and collaboration with labs in Berkeley, Leiden and Berlin.

In 1973, Stemler and Govindjee [5] worked out a procedure (by flushing isolated broken chloroplasts with nitrogen in a medium containing a high anion concentration at low pH) to remove bicarbonate; they obtained a large (5-fold) and reproducible effect of bicarbonate on the DCPIP Hill reaction in these samples; they suggested that HCO_3^- was bound in darkness and released in light. In view of their experiments with diphenylcarbazide (DPC), they had suggested that the effect was only on the O_2 -evolving side of PSII. This suggestion was challenged in 1975 by Wydrzynski and Govindjee [24] (see below).

In early 1974, Stemler and Govindjee [101] proceeded to perfect the methods of HCO_3^- -depletion further including the effects of light intensity and differences between the rates of O_2 evolution and ferricyanide reduction, again in broken maize chloroplasts, suggesting the possible existence of non- O_2 -evolving centers, and even an effect of bicarbonate on the rate of photoinactivation. These concepts still remain to be further investigated.

In 1974, Stemler and Govindjee [102] reported, working still with broken maize chloroplasts, complex effects of bicarbonate on Chl *a* variable fluorescence induction and delayed light emission, including an initial faster rise of Chl *a* fluorescence (from the minimum “O” level to the intermediate “I” level) in HCO_3^- -depleted conditions (in hindsight, a hint of an effect on the electron acceptor side); they suggested that HCO_3^- may stabilize the S_1 state in the dark, and, simply, bicarbonate is of critical importance in the initial photochemical process.

In collaboration with Gerald Babcock, then at the University of California, Berkeley, Stemler and Govindjee, in 1974 [23] presented the following findings: (1) bicarbonate decreases the probability of so-called “misses” in the system; (2) the turnover time of PSII is increased by $\text{CO}_2/\text{HCO}_3^-$ -depletion since the rate of dark relaxation of the S-states ($\text{S}_1' \rightarrow \text{S}_2$; $\text{S}_2' \rightarrow \text{S}_3$) is severely retarded in bicarbonate depleted broken maize chloroplasts (Fig. 4); in our current understanding, this may be either due to effects on the PSII acceptor or the donor or both sides; (3) the final O_2 -evolving reaction, after accumulation of four positive charges, is independent of bicarbonate; and (4) HCO_3^- has no effect on the dark deactivation of the higher oxidation states, S_2 and S_3 .

In 1975, Wydrzynski and Govindjee [24], as mentioned above, provided the first evidence that there was a clear effect of bicarbonate on the electron acceptor side of PSII: (1) absence of HCO_3^- led to a faster rise of Chl *a* fluorescence (reflecting reduction of Q_A to Q_A^-) in systems where the O_2 -evolving system was blocked (e.g., by Tris-washing) and artificial electron donors (e.g., NH_2OH , MnCl_2 , hydroquinone and even DPC) were added to replace water; (2) effect of increasing concentrations of the herbicide 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU; diuron), which blocks electron flow from Q_A^- to Q_B , mimics increasing $\text{CO}_2/\text{HCO}_3^-$ -depletion conditions (Fig. 5). These results leave no doubt about an effect of bicarbonate on the electron acceptor side of PSII. This, of course, does not mean that there is no bicarbonate effect on the donor side of PSII.

In 1976, Jursinic, also in Govindjee's lab, and in collaboration with Warden [103] demonstrated a major effect of bicarbonate on the electron acceptor side of PSII by using three separate and independent methods: EPR signal II “very fast”, corresponding to tyrosine Z radical, fast Chl *a* fluorescence yield changes; and delayed light emission; although no effects were observed on the electron donor side of PSII, a reversible inactivation of PSII RC activity was observed.

In 1976, Govindjee, in collaboration with Pulles, R. Govindjee, Van Gorkom and Duysens [104], discovered, using spinach chloroplasts, that HCO_3^- -depletion inhibits the re-oxidation of the reduced form of the secondary electron acceptor Q_B (Q_B^{2-}) by the PQ pool. Results on the effects of DCMU supported this conclusion. Flash-number dependent measurements on Chl *a* fluorescence yield established that in HCO_3^- -depleted samples, the “two-electron gate”, on the electron acceptor side of PSII, was non-functional (Fig. 6).

In 1977, Khanna et al. [105] performed the first, so-to-say, biochemical surgery of the bicarbonate effect, using artificial electron acceptors (silicomolybdate (SM), oxidized diaminodurene (DAD), and methyl viologen (MV)) and donors (DPC, reduced DAD), acting at specific sites, and the inhibitors (2,5-dibromo-3-methyl-6-isopropyl-*p*-benzoquinone (DBMIB) and DCMU) of electron flow, also at specific sites (see Fig. 7). Results were clear: (1) there was no bicarbonate effect on Photosystem I (PSI); (2) there was no bicarbonate effect on the water-oxidizing side of PSII; and (3) there was a definite inhibition of electron flow on the PSII acceptor side, in agreement with the work of Wydrzynski and Govindjee [24].

In 1977, in collaboration with Siggel and Renger, in Berlin, Khanna and Govindjee performed absorption spectroscopy to decipher the nature of electron carriers on the PSII acceptor side that were affected by $\text{CO}_2/\text{HCO}_3^-$ -depletion and the reversal after HCO_3^- re-addition [106]. The formation of Q_B^{2-} was reversibly slowed down, by a factor of 10–20 fold, from ~500 μs to ~8 ms. However, a much larger effect of $\text{CO}_2/\text{HCO}_3^-$ -depletion was in the slowing down of the reduction of PQ pool to ~100 ms; this was consistent with the measurements of Govindjee et al. [104], where the fluorescence decay after the 3rd and subsequent flashes was in the range of 150 ms; this was also reflected in the slowed reduction of oxidized P700. This is clearly the major bottleneck produced by bicarbonate depletion.

3.1.2.2. 1980s: further research at UIUC, and collaboration with other labs in Berlin and in Wako Shi (Japan).

In view of the fact that reduction of Q_B to PQH_2 requires protonation, it became obvious that bicarbonate must be playing a role through protonation, and, thus, in 1980, in collaboration with Junge's research group, in Berlin, Khanna et al. [107] measured the effect of HCO_3^- -depletion on the proton uptake and release, using pH indicator dyes neutral red (internal space) and bromo-cresol purple (external space); the results of HCO_3^- -depletion on protons were remarkable: not only was the release of protons into the internal space dramatically reduced, but there was no proton uptake by the PQ pool at the outer side of the membrane (Fig. 8). Whatever was the detailed mechanism, effects on protonation by bicarbonate depletion on PSII were firmly established.

In 1981, and in collaboration with the research group of Arntzen, and with Van Rensen, Khanna et al. [108] provided information suggesting that the binding of bicarbonate is on the same protein that binds the herbicide atrazine; further results suggested complete inactivation of a part of the total number of electron transport chains. These conclusions were based on: (1) a shift in the binding constant of atrazine in bicarbonate-depleted thylakoid membranes indicating decreased affinity of atrazine; (2) trypsin treatment, which modifies PSII at the level of Q_B , strongly diminished stimulation by bicarbonate addition to HCO_3^- -depleted thylakoids. These conclusions were confirmed by measurements on atrazine-resistant plants (Fig. 9).

In 1982, in collaboration with Van Rensen, Vermaas et al. [109] used the herbicide ioxynil, which is different from atrazine used earlier by Khanna et al. [108]; inhibition of electron transport by ioxynil increased at decreasing bicarbonate levels (Fig. 10). An interesting




Fig. 6. Increase of DCMU-induced Chl *a* fluorescence as a function of flash number in HCO₃⁻-depleted (closed squares), HCO₃⁻-depleted plus 20 mM NaHCO₃ (open squares), and control (open circles) spinach chloroplast suspensions as measured by Govindjee et al. [104]. Other conditions: [Chl] = 20 μg ml⁻¹; [DCMU] = 5 μM. The measurements were done in the presence of 1 mM NH₂OH used as an artificial electron donor. Modified and adapted from [104].

528 conclusion of this study was that the binding sites of herbicide and
 529 bicarbonate, although similar, are not identical.
 530 • In 1984, Blubaugh and Govindjee [30] came to the conclusion that
 531 bicarbonate has 2 binding sites: (1) a high affinity binding site
 532 close to where DCMU binds; this binding is inhibited by light; and
 533 (2) a low affinity binding site, which requires light, and is where
 534 bathocuproine may bind, and, thus, this could be the one effect on
 535 the donor side of PSII. These results and conclusions need further
 536 investigations. Considering the high resolution structure of PSII
 537 [17] where only one bicarbonate site has been seen, all experiments

dealing with two bicarbonate binding sites need to be re-examined
 and proven by additional experiments using newer methodologies.
 • In 1984, Eaton-Rye and Govindjee [110] extended the conclusions
 of Khanna et al. [105], using MV as electron acceptor, and provided
 additional evidence that there was no effect of bicarbonate in PSI,
 and that there was a specific effect on the PSII electron acceptor
 side (reduction of PQ) that was not dependent on the use of formate
 to remove bicarbonate. They suggested “the observed large slow
 component in HCO₃⁻-depleted samples results from an altered
 equilibrium of Q_A⁻ with PQ and/or PQH₂ at the Q_B binding site”.
 This conclusion was consistent with their hypothesis that removal
 of HCO₃⁻ results in a retardation of the PQ/PQH₂ exchange reactions
 of the two-electron gate. Further, it has been suggested that this
 may be due to changes in the association constants for one or
 more of the PQ/PQH₂ species and/or by affecting the protonation
 reactions of the partially reduced plastoquinone anion or the
 doubly reduced plastoquinol; it was only after the first full turnover
 of the two-electron gate that the full effect of HCO₃⁻ depletion could
 be observed. This also explained the observation of Govindjee et al.
 [104] and Robinson et al. [111] that the decay of Chl *a* fluorescence
 after the 1st flash is less inhibited than after the 3rd and subsequent
 actinic flashes, but intermediate after the 2nd flash.
 • In 1984, Govindjee et al. [112], working in the laboratory of Inoue
 (Wako Shi) in Japan, confirmed, through thermoluminescence (TL)
 measurements that the bicarbonate depletion affected PSII on the
 electron acceptor side, in the Q_AQ_B region. They discovered (1) a 6–10 °C
 shift, to a higher temperature, in the S₂Q_B⁻ TL band; (2) a reduction
 in TL intensity upon prolonged depletion of bicarbonate; and (3) elimi-
 nation, after the first few flashes, of the characteristic period four oscil-
 lations in TL intensity as a function of the flash number. On the other
 hand, addition of DCMU produced the same S₂Q_A⁻ TL band, at about
 +20 °C in both depleted and reconstituted samples. These results sug-
 gest (1) the initial effect of CO₂/HCO₃⁻ depletion is to increase the acti-
 vation energy for S₂(S₃)Q_B⁻ recombination; (2) with further depletion,
 the incidence of this recombination decreases and the cycling of the
 S₂Q_B⁻ and S₃Q_B⁻ recombination is inhibited through effects at the Q_B
 apo-protein. These bicarbonate depletion effects were fully reversible
 if HCO₃⁻ was added to HCO₃⁻-depleted samples (i.e., reconstituted sam-

Some inhibitors, artificial electron donors and electron acceptors of photosynthetic electron transport




Fig. 7. Isolation of the photosynthetic electron transfer chain into several segments (1, 2, and 3) by using artificial electron donors and acceptors in combination with specific inhibitors of electron carriers. Abbreviations: DPC, diphenylcarbazine; SM, silicomolybdate; BQ, benzoquinone; DAD, diaminodurene; DBMIB, 2,5-dibromo-3-methyl-6-isopropyl-p-benzoquinone; DCMU, diuron (3-(3,4-dichlorophenyl)-1,1-dimethylurea); MV, methyl viologen; Cyt *f*, cytochrome *f*; PC, plastocyanin; P700, RC Chl of PSI; A₀, and A₁, primary electron acceptors of PSI; FeS, iron sulfur centers of PSI. Other abbreviations are as in Fig. 3. Modified from [105].




Fig. 8. Flash-induced kinetics of proton release as measured by Khanna et al. [107] in control and CO_2 -depleted spinach thylakoids. (A) Proton release kinetics monitored by absorption changes of neutral red (NR) at 524 nm. Signals represent a difference between two transient signals (signal obtained in the absence of imidazole minus signal obtained in the presence of imidazole). The assays were performed in a medium containing 20 mM KCl, 2 mM MgCl_2 , 0.5 mM $\text{K}_3[\text{Fe}(\text{CN})_6]$, 0.3 μM nonactin, 10 μM NR, and 1.3 mg ml^{-1} bovine serum albumin (BSA) at pH 7.0. (B) Proton release kinetics as indicated by absorption changes of bromocresol purple (BCP) at 574 nm. Reaction medium (pH 6.4) contained 20 mM KCl, 2 mM MgCl_2 , 0.5 mM $\text{K}_3[\text{Fe}(\text{CN})_6]$, and 20 μM BCP. In both cases the final concentration of Chl in reaction mixture was 10 $\mu\text{g ml}^{-1}$. For illumination, saturating single-turnover flashes ($\tau_{1/2} = 15 \mu\text{s}$) were used, and the obtained signals were averaged over 10 flashes. Dark time between flashes was 10 s. Modified and adapted from [107].

576 ples). A conformational change of the PSII complex in the region of the
577 Q_B apo-protein was suggested to be responsible for these effects.
578 • The CO_2 concentration in water solutions ($[\text{CO}_{2(\text{aq})}]$) is a function of
579 Henry's Law of solubility and the partial pressure of CO_2 (g) in the
580 air above the water (see Fig. 1). Concentrations of other inorganic
581 carbon species, i.e., HCO_3^- and CO_3^{2-} , vary with pH, and therefore,
582 the ratio $[\text{HCO}_3^-]/[\text{CO}_{2(\text{aq})}]$ is pH dependent (for details, see [113]).
583 The total concentration of dissolved inorganic carbon increases at
584 the pH range between 6 and 9 due to an increase in HCO_3^- species.
585 In 1986, Blubaugh and Govindjee [114], taking advantage of the pH
586 dependence of the ratio $[\text{HCO}_3^-]/[\text{CO}_2]$ at equilibrium to vary effectively
587 the concentration of one species while holding the other species
588 constant, discovered that the Hill reaction was stimulated in
589 direct proportion with the equilibrium $[\text{HCO}_3^-]$, but was independent
590 of the equilibrium $[\text{CO}_2]$ (Fig. 11). Thus, they suggested that
591 HCO_3^- is the species, which binds to the effector site, while CO_2 is
592 the diffusing species [115].

- In 1988, Blubaugh and Govindjee [116], using kinetic analysis of rates
of electron flow versus $[\text{HCO}_3^-]$, came to the conclusion that there are
two high affinity bicarbonate binding sites, apparently with cooperative
binding. We now ask where is the second bicarbonate binding
site, if it really exists? On the PSII electron donor side? Or at another
site on the electron acceptor side? As mentioned above, since in a recent
high-resolution PSII structure there is no indication for two
 HCO_3^- molecules [17] (also see Section 3.2) the two binding site concept
needs to be re-examined with new experimental approaches.
Another conclusion was that bicarbonate is an essential activator for
PSII and that complete removal of HCO_3^- would result in zero electron
transport activity [116].
- In 1988, Eaton-Rye and Govindjee [117,118] provided a detailed
study of flash number dependent analysis of Chl *a* fluorescence
decay in spinach thylakoids at different pH values. The concept




Fig. 9. ^{14}C -labeled atrazine binding to CO_2 -depleted (open squares), CO_2 -depleted plus 20 mM NaHCO_3 (closed circles), and control (open circles) pea membrane thylakoids as reported by Khanna et al. [108]. The data were represented as plots of double reciprocal (mg Chl/nM bound atrazine) vs. $1/[\text{free atrazine}]$. Thylakoids were incubated at 23 $^\circ\text{C}$ with various concentrations of ^{14}C -labeled atrazine. The amount of bound atrazine was calculated from the difference between the total radioactivity added to the thylakoids and the amount of free atrazine found in the supernatant after centrifugation. Modified and adapted from [108].




Fig. 10. Double reciprocal plot of the ferricyanide Hill reaction rate (v_{Hill}) as a function of the bicarbonate concentration in the absence (closed circles) and presence (open circles) of 100 nM ioxynil in pea thylakoids as reported by Vermaas et al. [109]. The samples were incubated with bicarbonate for 2 min. The measurements of O_2 evolution rates were done in the presence of 0.5 mM $\text{K}_3[\text{Fe}(\text{CN})_6]$. Ioxynil was added 3.5 min prior to the measurements. Adapted and modified from [109].

608 that bicarbonate was involved in protonation was fully supported: a
609 model of bicarbonate acting as a proton donor to the protein disso-
610 sciable group believed to participate in the protonation of reduced
611 Q_B was discussed, as well as the possibility of HCO_3^- being a ligand
612 to the NHI in the Q_A-Fe-Q_B complex of the PSII RC. In addition, in
613 **1988**, (1) Cao and Govindjee [119] reported a bicarbonate effect in
614 a cyanobacterium *Synechocystis* sp. PCC 6803; and (2) Garab and
615 collaborators [120] provided evidence through TL measurements
616 that CO_2 does affect charge accumulation in intact leaves.

617 • In **1989**, (1) in collaboration with the Crofts' lab, Govindjee et al.
618 [121], using fast fluorescence changes, failed to observe any signifi-
619 cant effect of bicarbonate on electron donation from tyrosine Z (Y_Z)
620 to P680 or in the formation of $P680Q_A^-$; and (2) in collaboration
621 with the lab of Colin Wraight [122], a total absence of CO_2/HCO_3^- -
622 depletion effect was observed between the quinones both in chro-
623 matophores and RCs in the purple bacterium *Rhodobacter* (*R.*)
624 *sphaeroides*. This was followed, in **1992**, by the work of Wang (in
625 Wraight's Lab at Urbana, IL) and Cao (in Govindjee's lab) who, in collab-
626 oration with Oesterhelts' lab in Munich [123] asked if bicarbonate in
627 PSII is equivalent of Glu (M234 in *R. sphaeroides*) in bacterial RCs in
628 binding to the NHI. Michel and Deisenhofer [124] had earlier suggested
629 this notion. None of the mutants of M-234, where Glu was changed to
630 Val, Gln or Gly, showed any difference in the HCO_3^- -reversible formate
631 effect, confirming the absence of bicarbonate effect in these anoxygenic
632 photosynthetic bacteria.

633 3.1.2.3. 1990s: continued research at UIUC, and collaboration with other
634 labs in the USA and in labs around the World (Canada, China, Israel, Finland,
635 Switzerland, France, Germany, and The Netherlands).

636 • Following the lead of Khanna et al. [108] and Vermaas et al. [109] that
637 had suggested an overlap of binding sites of bicarbonate and herbicides
638 in higher plants, Govindjee, working in collaboration with Vernotte,
639 Peteri, Astier and Etienne, found, in **1990** [125] that the herbicide-
640 resistant mutants of the cyanobacterium *Synechocystis* sp. PCC 6714,
641 that are altered in specific amino acids in their D1 protein, show differ-
642 ential sensitivity to formate treatment. Yield of O_2 in a sequence of
643 flashes, Chl *a* fluorescence transients and Chl *a* fluorescence yield
644 decay after a flash revealed that the resistance of cells to formate treat-
645 ment was in the following (highest to lowest) order: [double
646 D1-mutant] A251V/F211S > [single D1-mutant] F211S > wild type >

[single D1-mutant] S264A. These results established the involvement
647 of the D1 protein in bicarbonate/formate binding, but gave no further
648 clue to the precise site of binding. From the PSII crystal structure [17],
649 these residues are rather close to Q_B ; changes in these residues may per-
650 turb the proper binding of Q_B , giving rise to indirect effects on the bind-
651 ing of bicarbonate/formate. 652

653 • In **1991**, using membrane-inlet mass spectrometry (MIMS) and a in-
654 frared gas analyzer, Govindjee in collaboration with Weger, Turpin,
655 Van Rensen, Devos and Snel, [53] showed that formate replaces
656 HCO_3^- from its binding site in PSII (see Fig. 12 and legend for experi-
657 mental details). Addition of 100 mM formate to spinach thylakoids re-
658 leased from $\sim 0.4 HCO_3^-/CO_2$ to $1.3 HCO_3^-/CO_2$, confirms the earlier idea
659 [12,125] that the bicarbonate effect occurs through the binding of
660 HCO_3^- to PSII, and that the addition of formate removes HCO_3^-/CO_2
661 from its binding site, leading to inhibition of electron flow. This did
662 not support the experiments and conclusions of Alan Stemler [126]. It
663 appears that about 1 HCO_3^- (at pH 6.5) is released by formate addition.
664 Further, in 1995, Oscar et al. [127] established the "bound-bicarbonate"
665 rather than the "inhibitory anion or the empty site" hypothesis of
666 Jursinic and Stemler [128] by showing CO_2 release under their experi-
667 mental conditions. 667

668 • Further evidence that the D1 protein was involved in the HCO_3^- effect
669 on PSII was obtained, in **1991**, by Govindjee et al. [129], using a
670 D1-L275F strain and several other mutants of *C. reinhardtii*, in collab-
671 oration with labs at University of Geneva, Switzerland. The L275F mu-
672 tant failed to show the HCO_3^- -reversible formate effect suggesting to
673 the authors that a significant change in formate (bicarbonate) binding
674 had occurred in helix V of the D1 protein near His involved in NHI
675 binding. Further, with the exception of the S264A mutant, which is
676 considerably more sensitive to formate than the wild type, five
677 other different [V219I, A251V, F255Y, G256D and cell-wall deficient
678 CW-15] mutants displayed a relatively similar response to formate
679 as wild type. Absence of a formate effect on a PSII-lacking mutant
680 seemed to confirm the sole involvement of PSII in the 'bicarbonate ef-
681 fect'. These results suggested that specific areas of the D1-protein are
682 more important than the others in formate/bicarbonate binding, but
683 they did not give precise clues. Lack of effect may not only be due to
684 the geometric organization of the structure, but may also be due to a
685 replacement with similar residues. The search continued. 685

686 • In **1991**, Xu and Govindjee [130], in collaboration with the laboratory
687 of Tony Crofts, presented a detailed kinetic investigation on spinach
688 thylakoids, as well as a model of HCO_3^- -reversible formate/formic
689 acid 688




Fig. 11. The rate of 2,6-dichlorophenolindophenol (DCPIP) reduction measured by Blubaugh and Govindjee [114] in CO_2 -depleted thylakoids as a function of the equilibrium CO_2 (A) and HCO_3^- (B) concentrations. The reduction rate of DCPIP was calculated from the decrease in absorbance at 600 nm and normalized to the control rate. The control rates (in μmol (DCPIP^{red}) mg (Chl)⁻¹ h^{-1}), estimated separately for each curve (pH value) by adding 2.5 mM HCO_3^- to the CO_2 -depleted samples, were the following: 209 at pH 6.31 (open squares); 212 at pH 6.54 (open diamonds); 191 at pH 6.67 (open circles); and 192 at pH 6.87 (open triangles). $NaHCO_3$ was added 3 min prior to illumination. Inset in (A): the effect of the equilibrium $[HCO_3^-]$ on the Hill reaction, with the $[CO_2]$ held constant at 0.1 mM. Inset in (B): the effect of the equilibrium $[CO_2]$ on the Hill reaction, with the $[HCO_3^-]$ held constant at 0.2 mM. Modified and adapted from [114].




Fig. 12. First detection of formate-induced release of CO₂ from spinach thylakoids as measured by MIMS by Govindjee et al. [53]. The addition of formate (to 100 mM) induced a rapid increase in the CO₂ signal (which corresponded to a formate injection artifact, i.e. formate blank) followed by a slow release of CO₂ from thylakoids (left). Repetitive addition of formate to the same sample induced only the initial rapid CO₂ release related to artifact of formate injection (right). CO₂ was continuously monitored at $m/z = 44$. The measurements were performed at 20 °C and pH 6.5 in the presence of external carbonic anhydrase (final concentration of 0.5 μg/ml) in order to facilitate the equilibration between inorganic carbon species.

Modified and adapted from [53].

acid effect. In agreement with earlier reports [104,118], electron flow from Q_A^- to Q_B^- was shown to be slowed down, and the notion that CO₂/HCO₃⁻-depletion leads to a loss of protonation necessary for stabilization of Q_B^- became the dominant mechanism. However, their conclusion that it is formic acid, not formate, which binds to the acceptor side of PSII has not been pursued since then, and it remains to be further investigated and deserves additional studies.

- In their detailed review in 1988, Blubaugh and Govindjee [12] had presented models for HCO₃⁻ action in PSII, and had also suggested that positively charged Arg residues may be involved in bicarbonate binding. In order to test this idea, Cao et al. [131], in 1991, made D2-R233Q and D2-R251S mutants in the cyanobacterium *Synechocystis* sp. PCC 6803, and, based on both O₂ evolution and Chl *a* fluorescence measurements, suggested that these two Arg residues must be somehow involved in bicarbonate binding in PSII. In the current high-resolution PSII structure [17], these two residues are located on the stromal surface 15–16 Å away from bicarbonate, and thus their effect may be indirect, through effects on the hydrogen-bonding network linking the bicarbonate to the stromal surface.
- In 1992, in collaboration with Pfister and Strasser's research group, Govindjee et al. [132] extended the earlier work on several herbicide resistant D1-C. *reinhardtii* mutants [129] and concluded that D1-S264, but not D1-L275, D1-F255 and D1-V219, plays an important role in the functioning of HCO₃⁻ and PQ in PSII; the role of D1-G256 could not be determined in this study. (See also [133] for the role of D1-S264 and the absence of the role of D1-L275.) The high-resolution structure of PSII [17] now shows that D1-S264 is hydrogen-bonded to Q_B^- (see Section 3.2).
- The role of various D1 amino acids in the HCO₃⁻ effect, this time, by using herbicide-resistant mutants of the cyanobacterium *Synechococcus*

sp. PCC 7942, was examined by Cao and Xiong, in 1992, in collaboration with Hirschberg and Ohad, in Israel [134]. Interestingly, the hierarchy of the equilibrium dissociation constant for bicarbonate (highest to lowest) was D1-F255L/S264A > D1-F255Y/S264A ~ D1-S264 ~ D1-F255Y > Wild type, establishing the importance of D1-S264 and D-F255 in the HCO₃⁻ binding niche directly or indirectly. Again, a role of bicarbonate in protonation and stabilization of Q_B^- was emphasized, a recurring concept since the earlier observations [107].

- By 1993, the following conclusions were made [135]: (1) formate, azide, nitrite and nitric oxide inhibited electron flow in thylakoids and cells, and these effects were significantly and uniquely reversed by bicarbonate; (2) with formate treatment, a remarkably strong HCO₃⁻-reversible slowing down of Q_A^- reoxidation after the second and subsequent flashes, but not after the first flash, was observed; (3) a hypothesis was in place suggesting that bicarbonate functions as a proton shuttle stabilizing the binding niche of Q_B^- and stimulating PQH₂ formation (and, perhaps, even its oxidation) in some manner; (4) this effect somehow involves both D1 and D2, directly, or indirectly, particularly the region where herbicides bind, and part of this was based on several mutant studies (e.g., D1-S264A, D1-L275F, D2-R251S, D2-R233Q, D2-R139H, among others); (5) possible involvement of "Fe" in the " Q_A -Fe- Q_B " complex was also implicated; (6) this effect was unique to PSII since electron transport in the " Q_A -Fe- Q_B " complex of both green and purple bacteria (including M-E234G, Q and V mutants) was insensitive to HCO₃⁻-reversible inhibitors.
- In 1995, Mäenpää et al. [136] made an interesting observation in *Synechocystis* sp. PCC 6803: HCO₃⁻-reversible formate effect on $Q_A Q_B$ was several fold less in the CA1 mutant (that had Glu 242, Glu 243, and Glu 244 deleted, and where Gln 241 was changed to His; these changes being in the de-loop of the D1 protein). These

results may be related to differences in the accessibility of the anions and/or due to changes in the redox properties of Q_A/Q_A^- in the mutant—perhaps, an indirect effect.

- In **1996**, in collaboration with the research group of Sayre, attempts were made to test the importance of D1-R269 in *C. reinhardtii* [137]; it was difficult to obtain firm conclusions since the used D1-R269G mutant was unable to grow photosynthetically and to evolve O_2 ; it had many defects. In the current high-resolution PSII structure [17], D1-R269 is hydrogen-bonded to D2-T243, which is probably needed to maintain the proper orientation of D2-Y244 in order for it to be able to hydrogen-bond to the bicarbonate (see Section 3.2).
- In **1996**, Xiong et al. [138] presented a 3-dimensional model of the D1/D2 protein and the cofactors, using the bacterial RCs, and predicted the HCO_3^- binding niche in PSII; it was modeled in the NHI site, providing a bidentate ligand to the iron. In their model, a bicarbonate ion that was suggested to be stabilized by D1-R257, was said to donate a proton to Q_B^{2-} through the D1-H252 residue, whereas a water molecule was proposed to donate another proton to Q_B^{2-} ; Xiong et al. also proposed a positively charged water channel, near Q_B and the NHI, for transporting water and HCO_3^- . It is now indeed known [17] that D1-H252 is hydrogen-bonded to Q_B through D1-S264, and there are water molecules close to D1-H252 that could serve as proton donor to Q_B (see Section 3.2). However, D1-R257 is at a distance of 8.6 Å from Q_B and further away from the bicarbonate; thus, it cannot be directly involved in this hydrogen-bond network.
- In **1997**, Govindjee et al. [7] presented data on Chl *a* fluorescence yield changes after light flashes 1–6 in spinach thylakoids at pH 6.0; they showed a bicarbonate effect on both the electron donor and electron acceptor sides in the same samples. The donor side effect was shown by a decrease in maximum fluorescence, and the acceptor side effect by a slowing down of the fluorescence decay due to Q_A^- oxidation. Using a sensitive differential infra-red gas analyzer they showed the presence of 0.8–1.25 bicarbonate ions bound per PSII RC in maize and pea thylakoids. These results were in agreement with earlier published data obtained by time-resolved MIMS on spinach thylakoids [53] (Fig. 12). Govindjee et al. [7] suggested that bicarbonate bound to the acceptor side is required for PSII activity, both on the acceptor and the donor sides in the same experiment and in the same sample; in this hypothesis, conformational changes may need to be invoked.

3.1.2.4. 2000s: new conclusions, collaboration with research group of Tony Crofts.

- In **2008**, Rose et al. [139], using both Chl *a* fluorescence, and TL measurements, provided the following conclusions on the D1-R257 mutation (D1-R257E, D1-R257M, and D1-257K): although the forward rate of electron transfer from Q_A to Q_B was little affected, the two-electron gate on the acceptor side of PSII was thermodynamically perturbed in the R257 mutants; this led to a decrease in the overall electron transfer rate from water to PQ. The effects on equilibrium constants of the two-electron gate are likely due to changes in coulombic fields on changing the net charge in the neighborhood of the Q_B site, suggesting that the electrostatic environment plays an important role in the mechanism of PSII. The bicarbonate-reversible formate effect on the Q_B site had been shown to be on the protonation events at this site [117,118]. Dramatic differences of the bicarbonate effect on the D1-R257 mutants, observed earlier [140], might thus have a basis in changes in the redox potential and the stability of the Q_B site, observed in this research. It, thus, seems that although D1-R257 is not close to the binding site of HCO_3^- on the NHI, it has a significant effect on the PSII reactions in the Q_B -region.
- As the model for the role of HCO_3^- had been evolving, it was generally thought that the first proton for the stabilization of Q_B^- came from D1-H252, and, thus, removal of HCO_3^- did not exhibit its major effect on the electron transport from the reduced Q_A to Q_B ,

but it had a large effect on the electron flow from the reduced Q_A to Q_B^- , and the succeeding reactions; the idea that bicarbonate provides this second proton, becoming carbonate, is the current picture. Carbonate, in turn, picks up a proton from D1-E244, finishing the cycle. The HCO_3^- ions (or water protons) outside the PSII complex provide the missing protons to the Glu (see current model in Section 3.2). The idea of involvement of D1-H252 in the first protonation was discussed by Petrouleas and Crofts [141], based on the experiments of Padden (see [142]; and paper in preparation). We note that depending upon the severity of HCO_3^- depletion procedure, an inhibition after the 1st flash is also observed explaining effects on TL band due to $S_2Q_B^-$ recombination (see earlier discussion).

3.1.3. Work around the World related to the site of bicarbonate binding

Research summarized below focuses on the studies related to key observations of bicarbonate binding to the electron acceptor side of PSII, mainly on the Q_A -NHI- Q_B niche and the PQ pool.

3.1.3.1. The 1980s.

- In **1984**, Vermaas and Rutherford [143] were among the first ones to focus on the relationship of bicarbonate to the Q_A -NHI- Q_B niche of PSII. They discovered that removal of HCO_3^-/CO_2 , in PSII membrane fragments from *Brassica napus*, led to a very large increase in the EPR signal at $g = 1.82$ that is due to the $Q_A^-Fe^{2+}$ complex, and, that this effect was fully reversible when bicarbonate was added back. This result identified bicarbonate to be either located near this complex, or, to play a crucial role in affecting the conformation of the Q_AFe complex.
- In **1987**, Diner and Petrouleas [144] showed reversible decrease in the quadrupole splitting of the NHI Mossbauer spectra, upon bicarbonate depletion. This confirmed the concept of bicarbonate acting on the electron acceptor side of PSII.
- In **1988**, Nugent et al. [145], using EPR measurements on both NHI ($g = 6$) and $Q_A^-Fe^{3+}$ ($g = 1.82$) in PSII particles, from both the thermophilic cyanobacterium *Phormidium laminosum* (Fig. 13A) and *Spinacea oleracea* (spinach) (Fig. 13B), suggested that bicarbonate binds close to the NHI and affects Q_A , Q_B as well as the NHI. Further, they found that the NHI was oxidized only when bicarbonate was present (also see [146]). These results supported the conclusions of Govindjee and coworkers (see Sections 3.1.2.1 and 3.1.2.2) that bicarbonate plays a central role in providing conditions for efficient electron flow on the acceptor side of PSII [11,147].
- In **1988**, Michel and Deisenhofer [124] in their perspective in the journal *Biochemistry* wrote "Having in mind the well-known effects of bicarbonate at the electron-accepting site of PSII, we consider bicarbonate as a likely candidate to be the fifth iron ligand in D1 and D2". They suggested that bicarbonate occupies the place of M-E232 of anoxygenic bacterial RC.

3.1.3.2. The 1990s.

- In **1990**, Diner and Petrouleas [25], using NO, instead of formate, to remove CO_2/HCO_3^- , showed that $g = 4$ EPR signal of $Fe^{2+}-NO$ was diminished when bicarbonate was added, favoring the concept that HCO_3^- is a ligand to the NHI.
- In **1991**, Diner et al. [148] presented a detailed overview on the iron-quinone electron acceptor complex of PSII. Here, they reviewed the literature on the bicarbonate effect in PSII and discussed various models for the binding and functioning of bicarbonate at the Q_A -NHI- Q_B complex.
- In **1995**, Hienerwadel and Berthomieu [149] provided the first IR spectroscopy evidence for bicarbonate binding on the acceptor side of PSII, using FT-IR difference spectroscopy, and ^{13}C -labeled HCO_3^- . Binding of bicarbonate to the NHI was strongly supported




Fig. 13. EPR data showing the effect of HCO_3^- removal on the acceptor side of PSII as reported by Nugent et al. [145]. (A) EPR spectra of the Q_A -NHI region ($g=1.82$) in dark-adapted PSII particles from the cyanobacterium *Phormidium laminosum* upon 5 min illumination at 77 K in the absence (spectrum 1) and the presence of 100 mM formate (spectrum 2). (B) EPR spectra of the NHI (Fe^{3+}) region ($g=6$) of the dark-adapted PSII particles from spinach in the absence (spectrum 1) and the presence (spectrum 2) of 100 mM formate. For further details see [145]. Modified and reproduced from [145].

by this study; it was suggested that bicarbonate is a monodentate ligand of the oxidized iron, but a bidentate ligand of the reduced form of iron, and exhibits hydrogen bonds with the protein.

3.1.3.3. The 2000s.

- In **2001**, Berthomieu and Hienerwadel [150] looked for the specific interactions of bicarbonate with the protein; here, they used lactate, glycolate and glyoxylate, instead of formate or NO, to remove inorganic carbon. Further, these authors concluded, from their studies, that one proton is released upon iron oxidation, and suggested that pH dependence of the iron couple may reflect deprotonation of D1-H215, a “putative” iron ligand located at the “ Q_B ” pocket. (This proton release was suggested to have a different mechanism from that involved in the functioning of bicarbonate.) They concluded that a ‘hydrogen network’ exists from the NHI towards the “ Q_B ” pocket involving bicarbonate and D1-H215 (see current model in Section 3.2).
- In **2008**, in search for proof (or absence of proof) for the binding of HCO_3^- to the electron donor side of PSII, Shevela et al. [41,151] re-examined and extended the MIMS experiments reported earlier by Govindjee et al. [53] and Stemler [126]. Govindjee et al. [53] had presented clear evidence for the release of $\text{CO}_2/\text{HCO}_3^-$ induced by formate addition (Fig. 12); however, the binding site for this anion was not specified in this study. Based on the previous experimental data, indicating the binding of HCO_3^- to the NHI at the acceptor side, it was assumed that formate removes HCO_3^- from this binding site. Formate, however, was reported to bind both at the acceptor and donor sides of PSII [152]. It was, therefore, unclear, from which binding side(s) in PSII the released CO_2 had originated in the previous study [53]. In the MIMS study of Shevela et al. some experiments were performed with an H_2^{18}O enrichment, which allowed the detection of CO_2 isotopologues at $m/z=46$ ($\text{C}^{16}\text{O}^{18}\text{O}$), and $m/z=48$ (C^{18}O_2). Since the Faraday cups used for the detection of $\text{C}^{16}\text{O}^{18}\text{O}$ and C^{18}O_2 were amplified by 10 and 100, respectively, than the one used for the detection of non-labeled CO_2 ($m/z=44$), the ^{18}O -enrichment greatly increased the sensitivity of the MS instrument (compare signal amplitudes in Fig. 14A, B, and C). The results obtained not only fully confirmed the formate-induced release of $\text{CO}_2/\text{HCO}_3^-$ reported earlier by Govindjee et al. [53] (Fig. 12), but also clearly demonstrated that the released $\text{HCO}_3^-/\text{CO}_2$ originates *only* from the acceptor side, and *not* from the donor side of PSII (for experimental details see Fig. 14 and its

legend). We also note here that, in the same year (**2008**), evidence for the binding of HCO_3^- on the electron acceptor side of PSII and the absence of bicarbonate bound to the donor side was presented in a FT-IR spectroscopy study by Aoyama et al. [42] and in a GC-MS study by Ulas et al. [55]. Thus, the focus of action on the mechanism remained on the Q_A -NHI- Q_B complex.

- In **2009**, Cox et al. [153] continued EPR studies on the Q_A -NHI- Q_B complex of PSII, initiated in the 1980s and 1990s (see above), but they added DFT calculations. They looked at the native $g\sim 1.9$ form as well as the $g\sim 1.84$ form, which is the well known signal in purple bacterial RCs (where bicarbonate does not bind, see Section 3.1.2.2) and that is occurring in PSII when they are treated with formate that removes $\text{CO}_2/\text{HCO}_3^-$. The calculations led Cox et al. to conclude that the doubly charged carbonate ion (CO_3^{2-}) is responsible for the $g\sim 1.9$ form of the semiquinone-iron signal; and carbonate, rather than bicarbonate (HCO_3^-), is the ligand to the NHI; the latter is in apparent contradiction to what we believe was the conclusion of Berthomieu and Hienerwadel (see above). It is highly likely that both bicarbonate and carbonate can bind to the NHI depending upon the precise physical and chemical status of the system since carbonate is formed from bicarbonate when the latter would be donating a proton to stabilize $\text{Q}_\text{B}^{\cdot -}$ (see Section 3.2)
- In **2009**, Takahashi et al. [154] dug deeply into the question of HCO_3^- binding at the NHI in PSII using FT-IR, as Berthomieu and Hienerwadel [150] had done, and included DFT calculations as well. Their study included specific ^{13}C -Tyr labeling together with a deuteration effect to provide evidence from Tyr IR modes to indicate Tyr involvement in hydrogen bonding to bicarbonate. The results obtained indicated that a Tyr (either D1-Y246 or D2-Y244; see Section 3.2) side chain in “a hydrogen bond donor-acceptor form” is strongly coupled to the NHI; this was suggested to provide a hydrogen bond to the oxygen of the bicarbonate ligand. Thus, Takahashi et al. were the first to propose that a key “Tyr residue coupled to the NHI may play a key role in the regulatory function of the iron-bicarbonate center by stabilizing the bicarbonate ligand and forming a rigid hydrogen bond network around the NHI.”
- In **2011**, Sedoud et al. [155] provided a thorough study on the effects of formate binding on the EPR of the quinone-NHI electron acceptor complex using light flash experiments and reached the conclusion that the effect was maximum after the 3rd flash indicating that the major effect of formate treatment ($\text{HCO}_3^-/\text{CO}_2$ removal) is on the $\text{Q}_\text{B}\text{H}_2$ exchange. This conclusion is in agreement with the earlier results of flash number dependence on Chl *a* fluorescence observed by Govindjee et al. [104] and on absorption changes by PQ, as measured by Siggel et al. [106]. However, this does not preclude, at all, the participation of bicarbonate in the protonation of $\text{Q}_\text{B}^{\cdot -}$. An integrated model would include both effects although the bottleneck reaction that would control the net electron flow may very well be this exchange reaction that would lead to slower oxidation of PQH_2 .
- In **2011**, Chernev et al. [156] investigated the NHI-(bi)carbonate complex using μs -resolution X-ray absorption spectroscopy (XAS) after laser flash excitation of PSII membrane particles. An interpretation of the observed spectral changes revealed that the coordination of bicarbonate at the Fe^{2+} may change from a bidentate to a monodentate ligation (carboxylate shift) after the formation of Q_A^- . Based on the obtained data and DFT calculations as well as on previous XAS experiments showing that no $\text{Fe}^{2+} \rightarrow \text{Fe}^{3+}$ transition occurs during the electron transfer from Q_A to Q_B in the type II photosynthetic RCs [157], Chernev et al. proposed that a coordination flexibility of the ligand (bicarbonate in PSII and glutamate in bacterial RCs) is essential for the functioning of the NHI-carboxyl complex in the interquinone electron transfer.
- In **2011**, Müh et al. [158] have beautifully reviewed PQ reduction in PSII. They suggest that one water molecule is there in the PSII structure that interacts with D1-H252, and two water molecules bridge




Fig. 14. Probing for binding sites of $\text{HCO}_3^-/\text{CO}_3^{2-}$ in PSII by isotope ratio MIMS in spinach PSII membrane fragments. These data confirmed the formate-induced release of CO_2 reported earlier by Govindjee et al. (see Fig. 12) and demonstrated that all released $\text{CO}_2/\text{HCO}_3^-$ originates from the electron acceptor and none from the donor side of PSII. (A) The addition of formate (to final concentration of 100 mM; black arrows) to PSII membranes at pH 6.3 and 20 °C induced a slow release of CO_2 (detected at $m/z = 44$) which was much above the artifact caused by injection of formate into the buffer with no samples (compare traces 1 and 2). Destruction of the possible binding site (the Mn_4CaO_5 cluster) via the addition of strong reductant NH_2OH (to final concentration of 7.5 mM; open arrows) does not lead to a release of $\text{CO}_2/\text{HCO}_3^-$ (compare traces 3 and 4). During the reduction NH_2OH is known to produce N_2O . In order to shift the signal of N_2O from $m/z = 44$ to $m/z = 46$, and thus to avoid possible overlay of the CO_2 and N_2O signals the ^{15}N -labeled NH_2OH was used for these experiments. (B) CO_2 release upon formate addition (to 100 mM) to 'control' PSII membranes (trace 1) is the same as in the case of PSII membranes without the Mn_4CaO_5 cluster (due to pre-incubation with 80 mM NH_2NH_2 for 75 min) (trace 2). CO_2 was detected at pH 6.3 and 20 °C as $^{16}\text{O}^{18}\text{O}$ at $m/z = 46$ due to ^{18}O -enrichment with H_2^{18}O (3%). (C) Formate-induced release of CO_2 (trace 1) compared with the absence of CO_2 release upon injection of NH_2OH (trace 2) as detected at $m/z = 48$ at pH 6.0 and 20 °C. To get the highest possible sensitivity the experiments were performed with high ^{18}O -enrichment level (~65%). All measurements were done in the presence of externally added carbonic anhydrase (to a final concentration of 3 $\mu\text{g}/\text{ml}$) to facilitate equilibration between CO_2 and HCO_3^- and by this to allow the detection of all dissolved inorganic carbon as CO_2 along. Modified and adapted from [41].

Shen reported, in 2003 [161], the PSII structure from *T. vulcanus* at a 3.7 Å resolution. These structures did not allow the assignment of bicarbonate in PSII, either on the donor, or the acceptor side. The first assignment of bicarbonate was reported by Ferreira et al., in 2004 [51], in their PSII structure from *T. elongatus* at 3.5 Å resolution, in which they assigned two HCO_3^- ions, one at the donor side and the other at the acceptor side. The bicarbonate at the donor side was assigned to be a direct ligand to the Mn_4CaO_5 cluster. The density that was assigned to a putative bicarbonate, however, was not found in the subsequent structures at higher resolutions of 2.9–3.0 Å [52,162]. In the most recent structure of PSII determined at a resolution of 1.9 Å [17], bicarbonate was also not found at the donor side. Since in this high resolution structure, all of the ligands for the 4 Mn ions and the Ca ion were determined, which showed that each of the Mn has 6 ligands and the Ca ion has 7 ligands, there is no room for the presence of a bicarbonate in the immediate ligand sphere of the Mn_4CaO_5 cluster, at least in the assembled, active PSII complex. It is also highly unlikely that a well-defined HCO_3^- could be missed in an electron density map with a resolution beyond 2.0 Å, as the electron density for the bicarbonate at the acceptor was clearly defined and visible [17]. One can assume that bicarbonate strongly bound to the Mn_4CaO_5 cluster might be lost due to reduction of high-valence Mn ions ($\text{Mn}^{\text{III}}\text{Mn}^{\text{IV}}_2$) to Mn(II), which is known to take place under X-ray doses used for structure determination by X-ray crystallography [163]. However, in view of recent MS and FT-IR data [41,42,55,151] showing the absence of tightly bound bicarbonate to the Mn_4CaO_5 cluster, this option can be excluded.

Fig. 15A shows the position of the bicarbonate on the acceptor side in a PSII monomer determined at 1.9 Å resolution [17]. While the global position of the bicarbonate could be assigned in the structures with a resolution in the range of 3.0–3.5 Å, its detailed environment including the presence and the positions of water molecules surrounding it has to be determined at a much higher resolution, which is now achieved at 1.9 Å. Based on this structure, the bicarbonate serves as a bidentate ligand to the NHI, which is located just under the surface of the stromal side of the membrane region. This bicarbonate is surrounded by hydrophilic residues and water molecules, indicating that it is in a highly hydrophilic environment.

As we can see from Fig. 15B, there is a very small proteinaceous region from the HCO_3^- toward the stromal solution; thus, protons from the stromal side are expected to have easy access to the site of bicarbonate. In order for an efficient and uni-directional transfer of protons to be able to occur, however, hydrogen-bond networks are expected to be present. In fact, well-defined hydrogen-bond networks have been found linking the bicarbonate to the stromal bulk solution. As shown in Fig. 16, the 3rd oxygen in the bicarbonate that was not ligated to the NHI is hydrogen-bonded to a water molecule (W1138A in the 1.9 Å structure, PDB ID: 3ARC). This water molecule has four hydrogen bonds with its neighboring groups, among which, two are Tyr and Ser residues of the D1 protein (D1-Y246 and D1-S268), and the 3rd one is another water molecule (W675A). This 2nd water molecule (W675A) extends the hydrogen-bond network to the stromal surface through another water molecule W2195D. A plausible hypothesis is: after reduction of Q_B by the reduced Q_A , protons could be easily taken in from the stromal bulk solution through this hydrogen-bond network, and transferred to the site of bicarbonate, which may be further transferred to the reduced Q_B through D1-H272 and D1-H215.

Both the 1st (W1138A) and the 2nd (W675A) water molecules, which are hydrogen-bonded to the bicarbonate, have a tetragonal configuration, bearing 4 hydrogen bonds with their neighboring molecules. The amino acid residues surrounding them thus seem to be important for holding these two water molecules in a proper position, in order to form the proper hydrogen-bond network connecting the bicarbonate to the stromal side. These residues include D1-S268, D1-Y246 for the 1st water molecule, and D1-E244, D2-T243 for the 2nd water molecule. Changes in one of these residues may therefore disturb the positions of the water molecules, and thereby disrupt the proper hydrogen-bond

D1-E244 and D1-Y246, and these could very well be involved in proton pathways (see Figs. 4A and 6A in [158]). They independently propose, as Govindjee, in collaboration with Crofts and Padden [142,159] has suggested that the first proton enters via D1-H252 and the second via D1-Y246, and that it may involve D1-E244.

3.2. The crystal structure at a resolution 1.9 Å and the current model for bicarbonate function

The crystal structure of PSII was first reported by Zouni et al. in 2001 [160] at a resolution of 3.8 Å from a thermophilic cyanobacterium *Thermosynechococcus* (*T.*) *elongatus*. Subsequently, Kamiya and




Fig. 15. (A) Structure of a PSII monomer determined at a 1.9 Å resolution [17]. View from the direction perpendicular to the membrane normal. Dashed lines represent the cytoplasmic (stromal) and luminal surface of the membrane, respectively. Color codes: green, D1; yellow, D2; cyan, CP47; dark pink, CP43; red, PsbL; light pink, PsbO; light blue, PsbU. The blue ball in the middle represents the NHI, and a magenta molecule above the NHI represents bicarbonate. (B) An enlarged view of the NHI and bicarbonate region shown in (A). The color codes are the same as for (A).

networks. D2-K264 is not hydrogen-bonded to any of the two water molecules, but is hydrogen-bonded to D2-E242, and also close to D1-E244, one of the residues hydrogen-bonded to the 2nd water molecule. Alteration of D2-K264 may therefore perturb the orientation of D2-E242, resulting in an effect on the position of the 2nd water molecule.

We know that there is another short H-bond network that connects Q_B to the stromal surface, which is composed of D1-S264 and D1-H252. D1-H252 is located in a small dent in the stromal surface and is hydrogen-bonded to a water molecule directly, which is further hydrogen-bonded to another water molecule. A number of additional water molecules are found in the vicinity of these water molecules, indicating that D1-H252 is located in a highly hydrophilic area. Thus, protons may also be easily taken from this area of the stromal surface and transferred to Q_B through D1-H252 and D1-S264. In view of the previous functional studies [142], it is plausible to suggest that the first proton to protonate Q_B^- is taken up through D1-H252 and D1-S264, and the second proton is transferred via bicarbonate to D1-H272 and D1-H215, and, finally to Q_B through the H-bond network (see Fig. 16, and its legend). HCO_3^- that must become CO_3^{2-} ,

after giving up its proton to Q_B^{2-} , may get its proton back from the stroma via D1-E244 [158,159]. However, further functional studies are required to prove or disprove the order of these protonation events. In addition, since there are indications for a change of the bicarbonate coordination to the NHI from bidentate to monodentate upon electron transport from Q_A towards Q_B [150,156], there might be alternative proton paths newly created by possible accompanying conformational changes.

In view of the above picture of the environment and plausible function of bicarbonate, we recommend comparative biochemical and biophysical studies on appropriate site-directed mutants of D1-E244; D1-Y246; D1-S268; D2-T243; D2-E242; and D2-K264.

4. Uniqueness of role of bicarbonate in oxygenic photosynthesis

The requirement of PSII for bicarbonate (carbonate) has been observed at the level of intact leaves, isolated thylakoids and PSII-enriched membrane fragments from plants, algae, and cyanobacteria, but never in the RCs of anoxygenic photosynthetic bacteria (see sections above and references therein). It appears, therefore, that by being a ligand to the NHI between Q_A and Q_B , and binding to amino acids of the D1 and D2 proteins of PSII in these organisms, bicarbonate/carbonate plays a unique role only in oxygenic photosynthesis: it stabilizes the Q_A -NHI- Q_B structure of the PSII RC, and, thus, allows efficient electron transport and protonation of Q_B^- via certain amino acids around Q_B (Fig. 16). We ask: why does the PSII RC have, unlike its bacterial cousin, a bicarbonate ion liganded to its NHI? The simple answer is that it may have a regulatory function here in PSII electron flow. Under normal conditions, bicarbonate may be bound and function in protonation events, as discussed above; however, when the plant is exposed to drought, high light and high temperature, the stomata may close, leading to a decrease of the internal $[CO_2]$. Similar decreases in $[CO_2]$ are expected in algae and cyanobacteria that do not have stomata [164]. This would lead to a decrease in $[HCO_3^-]$ limiting PSII activity.

The effect of bicarbonate depletion within PSII is not only on the electron acceptor side, but also on the donor side, although the exact location (or binding site) responsible for this effect of HCO_3^- is not known [9,14,20]. The effect of HCO_3^- on the water-oxidizing side of PSII has also been seen *in vitro* in all oxygenic organisms (higher plants, algae, and cyanobacteria) (for details see Section 2). There is, however, lack of observations of this effect in intact organisms. Many experimental data obtained on isolated PSII membrane fragments and PSII core preparations are consistent with a unique role of HCO_3^- in initiating and/or facilitating assembly of the inorganic core of the OEC from OEC-depleted PSII RCs (e.g., arising as a result of disassembly of the OEC under stress conditions or when newly synthesized) and Mn^{2+} ions (reviewed in [61]). There are also indications for the functioning of HCO_3^- in the assembled OEC [14]. Thus, for instance, newer data suggest that mobile (loosely bound or even non-bound) bicarbonate may facilitate deprotonation of the Mn_4CaO_5 cluster (opposite to the protonation reactions assigned for the ‘acceptor-side’ HCO_3^-) [32,34]. By “picking up” the protons that are produced during water splitting, HCO_3^- *per se* or in concert with CA may play a regulatory function against over-acidification of the lumen in the proximity of the water-oxidizing site, and by this, protect the OEC against destabilization and predisposition to photoinhibition.

5. Bicarbonate and evolutionary development of the O_2 -evolving Photosystem II

All O_2 -producing photosynthetic organisms (cyanobacteria, green algae, and plants) have the same Mn_4CaO_5 inorganic core and very similar RC core proteins forming the basis for PSII capable of catalyzing oxidation of water. The available geological and geochemical data indicate that nature created this single type of enzyme as early as 3.2 Ga or as late as 2.4 Ga ago [165–168]. The role of bicarbonate (CO_2) in the evolutionary




Fig. 16. Hydrogen-bond networks around Q_B , the NHI and bicarbonate. Blue lines represent coordination bonds, and dashed lines in cyan indicate hydrogen-bonds. Arrows in dashed, black lines indicate possible flow of protons towards the Q_B molecule. One of the protons (say the first one) may be picked up through D1-H252 and D1-S264 to protonate Q_B^- , while the second one may be transferred through bicarbonate, D1-H272, and D1-H215 to Q_B (see Section 3.2 for more details). Figure based on data of Umena et al. [17].

1139 development of the first O_2 -evolving cyanobacteria-like organisms is obvious, since the presence of HCO_3^-/CO_3^{2-} bound between Q_A and Q_B in the
 1140 RC is unique, as it exists only in oxygenic photoautotrophs, whereas it is
 1141 absent in all anoxygenic photosynthesizers [122,123]. The coupling of bicarbonate as a ligand to facilitate Q_B^- protonation and, thus, the electron
 1142 transfer in the first O_2 -producing organisms *via* replacement of the Glu ligand in anoxygenic bacterial RCs (as first suggested in [124]; for further
 1143 details, see Section 3.1.3.1) could be simply an additional evolutionary step from anoxygenic towards oxygenic photosynthesis [169].

1144 There are also indications for a key role of Mn–bicarbonate complexes in the evolutionary origin of the water-oxidizing inorganic core of the
 1145 OEC of PSII [59,170]. The unique capability of bicarbonate to form easily oxidizable complexes with Mn ions has been demonstrated in numerous
 1146 electrochemical and EPR studies (see, for instance, [62,64,65]). Since the oxidation potentials of the Mn^{2+} – HCO_3^- complex (520–680 mV) were
 1147 found to have close values to the midpoint redox potentials of the primary electron donor (P) in the RCs of non-oxygenic bacteria, Dismukes et al.
 1148 [170] suggested, that these complexes (which could be formed under much higher concentrations of dissolved CO_2 (HCO_3^-) in the ancient
 1149 ocean than at present) were probably used as a source of electrons by some Archean anoxygenic bacteria—ancestors to the first oxygenic cyanobacteria. Results obtained both on wild type contemporary purple
 1150 bacteria [171] and on mutants of *R. sphaeroides* with modified midpoint redox potentials of the P/P⁺ RC couple [33] support this idea by showing
 1151 that the formation of the Mn^{2+} –bicarbonate complexes stimulate electron donation from Mn^{2+} to type II RCs of these anoxygenic
 1152 bacteria.

1166 6. Concluding remarks

1167 As proved by recent X-ray crystallography studies of PSII [16,17], in the cyanobacterial RC, there is only evidence for a single bound bicarbonate at the NHI. The evidence for “bicarbonate” as ligand to the
 1168 quinone–iron complex derived from a large body of data makes it clear that there is a role for this ligand *in vivo*. The presence of bicarbonate as a bidentate ligand to the NHI bridging Q_A and Q_B is now
 1169

1170 firmly established [16,17]. This set the key stone to a huge body of studies that have established a role of bicarbonate in facilitating proton
 1171 transfer and, thereby, accelerating electron transfer between Q_A , Q_B and from Q_B into the PQ-pool; the absence of bicarbonate might
 1172 down-regulate this electron transfer step. Since this action prevails in all oxygenic organisms, the structural and functional role of bicarbonate has arisen very early in evolution. There is also an effect on the
 1173 reoxidation of PQH₂. Comparative biochemical and biophysical studies on site-directed mutants of tyrosines near the HCO_3^- binding site is expected to provide key information on the mechanistic role of bicarbonate in these reactions.

1184 There is another, though less well defined role of bicarbonate on the donor side of PSII. A particular binding site close to the Mn_4CaO_5 cluster is absent in the high-resolution structure [17]. Since the roles of HCO_3^-
 1185 as a mobile substrate of PSII or as a direct tightly bound ligand to the Mn_4CaO_5 cluster are excluded by numerous studies, a direct involvement of HCO_3^-
 1186 in the water-oxidizing process can now be ruled out. There is, however, undeniable evidence for an essential role of HCO_3^- in the process of photoactivation. Further experiments are required to evaluate
 1187 the possibility of HCO_3^- involvement in the deprotonation reactions of the OEC. The indirect effects of HCO_3^- on water oxidation (such as, protection against thermoinactivation, photoinhibition, protein extraction, and
 1188 treatments with some reductants) need to be studied and characterized further. Moreover, one should clarify whether HCO_3^- ions have the same function on the donor side of PSII in intact photosynthetic systems.

1198 Acknowledgements

1199 DS acknowledges the financial support by the Norges forskningsråd. JJER was supported by a New Zealand Marsden Contract 08-UOO-043. Work in the JRS group was supported by grants from the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Yamada Science Foundation and the Mitsubishi Foundation. G is thankful to the Department of Plant Biology and the Office of Jeff Haas, at the University of Illinois at Urbana-Champaign, USA, for their support. During the final phase of this review, Govindjee was supported, by a visiting

1207 professorship, at the School of Life Sciences, Jawaharlal Nehru University, New Delhi, India. Lastly, our thanks go to the four reviewers for their
 1208 important suggestions that have led to a significant improvement in this
 1209 review.
 1210

1211 References

- 1212 [1] V.M. Krishnamurthy, G.K. Kaufman, A.R. Urbach, I. Gitlin, K.L. Gudiksen, D.B.
 1213 Weibel, G.M. Whitesides, Carbonic anhydrase as a model for biophysical and
 1214 physical–organic studies of proteins and protein–ligand binding, *Chem. Rev.*
 1215 **108** (2008) 946–1051.
- 1216 [2] T. Wydrzynski, K. Satoh, Photosystem II. The light-driven water:plastoquinone
 1217 oxidoreductase, in: Govindjee (Ed.), *Advances in Photosynthesis and Respiration*,
 1218 Springer, Dordrecht, 2005.
- 1219 [3] W. Martin, R. Scheibe, C. Schnarrenberger, The Calvin cycle and its regulation, in:
 1220 R.C. Leegood, T.D. Shakey, S. von Caemmerer (Eds.), *Photosynthesis: Physiology
 1221 and Metabolism*, Kluwer Academic Publishers, Dordrecht, 2000, pp. 9–51.
- 1222 [4] O. Warburg, G. Krippahl, Hill-Reaktionen, *Z. Naturforsch. B* **13** (1958) 509–514.
- 1223 [5] A. Stemler, Govindjee, Bicarbonate ion as a critical factor in photosynthetic oxy-
 1224 gen evolution, *Plant Physiol.* **52** (1973) 119–123.
- 1225 [6] O. Warburg, Prefatory chapter, *Annu. Rev. Biochem.* **33** (1964) 1–18.
- 1226 [7] Govindjee, C. Xu, J.J.S. van Rensen, On the requirement of bound bicarbonate for
 1227 photosystem II activity, *Z. Naturforsch.* **52** (1997) 24–32.
- 1228 [8] J.J.S. Van Rensen, Role of bicarbonate at the acceptor side of photosystem II,
 1229 *Photosynth. Res.* **73** (2002) 185–192.
- 1230 [9] A.J. Stemler, The bicarbonate effect, oxygen evolution, and the shadow of Otto
 1231 Warburg, *Photosynth. Res.* **73** (2002) 177–183.
- 1232 [10] V.V. Klimov, S.V. Baranov, Bicarbonate requirement for the water-oxidizing
 1233 complex of photosystem II, *Biochim. Biophys. Acta* **1503** (2001) 187–196.
- 1234 [11] Govindjee, J.J.S. Van Rensen, Bicarbonate effects on the electron flow in isolated
 1235 broken chloroplasts, *Biochim. Biophys. Acta* **505** (1978) 183–213.
- 1236 [12] D.J. Blubaugh, Govindjee, The molecular mechanism of the bicarbonate effect at the
 1237 plastoquinone reductase site of photosynthesis, *Photosynth. Res.* **19** (1988) 85–128.
- 1238 [13] Govindjee, J.J.S. Van Rensen, Photosystem II reaction centers and bicarbonate, in:
 1239 J. Deisenhofer, J.R. Norris (Eds.), *Photosynthetic Reaction Centers*, Academic
 1240 Press, Orlando, 1993, pp. 357–389.
- 1241 [14] J.J.S. Van Rensen, V.V. Klimov, Bicarbonate interactions, in: T. Wydrzynski, K. Satoh
 1242 (Eds.), *Photosystem II. The Light-Driven Water:Plastoquinone Oxidoreductase*,
 1243 Springer, Dordrecht, 2005, pp. 329–346.
- 1244 [15] Govindjee, J. Kern, J. Messinger, J. Whitmarsh, Photosystem II, *Encyclopedia of
 1245 Life Sciences (ELS)*, John Wiley & Sons, Ltd., Chichester, 2010, pp. 1–15.
- 1246 [16] A. Guskov, A. Gabdulkhakov, M. Broser, C. Glöckner, J. Hellmich, J. Kern, M. Frank,
 1247 W. Saenger, A. Zouni, Recent progress in the crystallographic studies of photosys-
 1248 tem II, *Chemphyschem* **11** (2010) 1160–1171.
- 1249 [17] Y. Umena, K. Kawakami, J.-R. Shen, N. Kamiya, Crystal structure of oxygen-
 1250 evolving photosystem II at a resolution of 1.9 Å, *Nature* **473** (2011) 55–60.
- 1251 [18] G. Renger, A.R. Holzwarth, Primary electron transfer, in: T.J. Wydrzynski, K. Satoh
 1252 (Eds.), *Photosystem II. The Light-Driven Water:Plastoquinone Oxidoreductase*,
 1253 Springer, Dordrecht, 2005, pp. 139–175.
- 1254 [19] G. Renger, Photosynthetic water splitting: apparatus and mechanism, in:
 1255 J.J. Eaton-Rye, B.C. Tripathy, T.D. Sharkey (Eds.), *Photosynthesis: Plastid Biology, En-
 1256 ergy Conversion and Carbon Assimilation*, Springer, Dordrecht, 2012, pp. 359–411.
- 1257 [20] I.L. McConnell, J.J. Eaton-Rye, J.J.S. Van Rensen, Regulation of photosystem II
 1258 electron transport by bicarbonate, in: J.J. Eaton-Rye, B.C. Tripathy, T.D. Sharkey
 1259 (Eds.), *Photosynthesis: Plastid Biology, Energy Conversion and Carbon Assimila-
 1260 tion*, Springer, Dordrecht, 2012, pp. 475–500.
- 1261 [21] A. Stemler, The functional role of bicarbonate in photosynthetic light reaction II, in:
 1262 Govindjee (Ed.), *Photosynthesis*, Academic Press, New York, 1982, pp. 513–538.
- 1263 [22] A. Stemler, Bicarbonate and photosynthetic oxygen evolution: an unwelcome
 1264 legacy of Otto Warburg, *Ind. J. Exp. Biol.* **36** (1998) 841–848.
- 1265 [23] A. Stemler, G.T. Babcock, Govindjee, Effect of bicarbonate on photosynthetic ox-
 1266 ygen evolution in flashing light in chloroplast fragments, *Proc. Natl. Acad. Sci. U.
 1267 S. A.* **71** (1974) 4679–4683.
- 1268 [24] T. Wydrzynski, Govindjee, New site of bicarbonate effect in photosystem II of
 1269 photosynthesis – evidence from chlorophyll fluorescence transients in
 1270 spinach-chloroplasts, *Biochim. Biophys. Acta* **387** (1975) 403–408.
- 1271 [25] B.A. Diner, V. Petrouleas, Formation by NO of nitrosyl adducts of redox components
 1272 of the photosystem II reaction center. 2. Evidence that HCO₃⁻/CO₂ binds to the
 1273 acceptor-side non-heme iron, *Biochim. Biophys. Acta* **1015** (1990) 141–149.
- 1274 [26] A. Stemler, Inhibition of photosystem II by formate. Possible evidence for a di-
 1275 rect role of bicarbonate in photosynthetic oxygen evolution, *Biochim. Biophys.
 1276 Acta* **593** (1980) 103–112.
- 1277 [27] H. Metzner, *Photosynthetic Oxygen Evolution*, Academic Press, London, 1978.
- 1278 [28] W. Kreuz, Considerations on water-splitting in photosynthesis, in: K. Colbow
 1279 (Ed.), *On the Physics of Biological Membranes*, Department of Physics, Simon
 1280 Fraser University, Vancouver, 1974, pp. 419–429.
- 1281 [29] D. Mende, W. Wiessner, Bicarbonate *in vivo* requirement of photosystem II in
 1282 the green alga *Chlamydomonas reinhardtii*, *J. Plant Physiol.* **118** (1985) 259–266.
- 1283 [30] D.J. Blubaugh, Govindjee, Comparison of bicarbonate effects on the variable
 1284 chlorophyll *a* fluorescence of CO₂-depleted and non CO₂-depleted thylakoids
 1285 in the presence of diuron, *Z. Naturforsch. C* **39** (1984) 378–381.
- 1286 [31] S.V. Baranov, A.M. Tyryshkin, D. Katz, G.C. Dismukes, G.M. Ananyev, V.V. Klimov,
 1287 Bicarbonate is a native cofactor for assembly of the manganese cluster of the
 photosynthetic water oxidizing complex. Kinetics of reconstitution of O₂ evolu-
 tion by photoactivation, *Biochemistry* **43** (2004) 2070–2079.
- [32] T. Shutova, H. Kenneweg, J. Buchta, J. Nikitina, V. Terentyev, S. Chernyshov,
 B. Andersson, S.I. Allakhverdiev, V.V. Klimov, H. Dau, W. Junge, G. Samuelsson, The
 photosystem II-associated Cah3 in *Chlamydomonas* enhances the O₂ evolution rate
 by proton removal, *EMBO J.* **27** (2008) 782–791.
- [33] L. Kalman, J.C. Williams, J.P. Allen, Energetics for oxidation of a bound manganese
 cofactor in modified bacterial reaction centers, *Biochemistry* **50** (2011) 3310–3320.
- [34] G. Ananyev, T. Nguyen, C. Putnam-Evans, G.C. Dismukes, Mutagenesis of CP43-
 arginine-357 to serine reveals new evidence for (bi)carbonate functioning in the
 water oxidizing complex of photosystem II, *Photochem. Photobiol. Sci.* **4** (2005)
 991–998.
- [35] G. Ulas, G.W. Brudvig, Zwitterion modulation of O₂-evolving activity of cyano-
 bacterial photosystem II, *Biochemistry* **49** (2010) 8220–8227.
- [36] P.A. Castelfranco, Y.K. Lu, A.J. Stemler, Hypothesis: the peroxydicarbonic acid
 cycle in photosynthetic oxygen evolution, *Photosynth. Res.* **94** (2007) 235–246.
- [37] R. Radmer, O. Ollinger, Isotopic composition of photosynthetic O₂ flash yields in
 the presence of H₂¹⁸O and HCl¹⁸O₂, *FEBS Lett.* **110** (1980) 57–61.
- [38] J. Clausen, K. Beckmann, W. Junge, J. Messinger, Evidence that bicarbonate is not the
 substrate in photosynthetic oxygen evolution, *Plant Physiol.* **139** (2005) 1444–1450.
- [39] W. Hillier, I. McConnell, M.R. Badger, A. Boussa, V.V. Klimov, G.C. Dismukes,
 T. Wydrzynski, Quantitative assessment of intrinsic carbonic anhydrase activity
 and the capacity for bicarbonate oxidation in photosystem II, *Biochemistry* **45**
 (2006) 2094–2102.
- [40] R. Radmer, O. Ollinger, Do the higher oxidation states of the photosynthetic O₂
 evolving system contain bound water? *FEBS Lett.* **195** (1986) 285–289.
- [41] D. Shevela, J.H. Su, V. Klimov, J. Messinger, Hydrogencarbonate is not a tightly
 bound constituent of the water-oxidizing complex in photosystem II, *Biochim.
 Biophys. Acta* **1777** (2008) 532–539.
- [42] C. Aoyama, H. Suzuki, M. Sugiura, T. Noguchi, Flash-induced FTIR difference spectro-
 scopy shows no evidence for the structural coupling of bicarbonate to the oxygen-
 evolving Mn cluster in photosystem II, *Biochemistry* **47** (2008) 2760–2765.
- [43] V.V. Klimov, S.I. Allakhverdiev, Y.M. Feyziev, S.V. Baranov, Bicarbonate require-
 ment for the donor side of photosystem II, *FEBS Lett.* **363** (1995) 251–255.
- [44] S.I. Allakhverdiev, I. Yruela, R. Picorel, V.V. Klimov, Bicarbonate is an essential
 constituent of the water-oxidizing complex of photosystem II, *Proc. Natl. Acad.
 Sci. U. S. A.* **94** (1997) 5050–5054.
- [45] R.J. Hulsebosch, S.I. Allakhverdiev, V.V. Klimov, R. Picorel, A.J. Hoff, Effect of bi-
 carbonate on the S₂ multiline EPR signal of the oxygen-evolving complex in
 photosystem II membrane fragments, *FEBS Lett.* **424** (1998) 146–148.
- [46] D.N. Shevela, A.A. Khorobrykh, V.V. Klimov, Effect of bicarbonate on the water-
 oxidizing complex of photosystem II in the super-reduced S-states, *Biochim.
 Biophys. Acta* **1757** (2006) 253–261.
- [47] I. Yruela, S.I. Allakhverdiev, J.V. Ibarra, V.V. Klimov, Bicarbonate binding to the
 water-oxidizing complex in the photosystem II. A Fourier transform infrared
 spectroscopy study, *FEBS Lett.* **425** (1998) 396–400.
- [48] V.V. Klimov, S.I. Allakhverdiev, S.V. Baranov, Y.M. Feyziev, Effects of bicarbonate and
 formate on the donor side of photosystem 2, *Photosynth. Res.* **46** (1995) 219–225.
- [49] V.V. Klimov, S.I. Allakhverdiev, Y. Nishiyama, A.A. Khorobrykh, N. Murata, Stabi-
 lization of the oxygen-evolving complex of photosystem II by bicarbonate and
 glycinebetaine in thylakoid and subthylakoid preparations, *Funct. Plant Biol.*
30 (2003) 797–803.
- [50] V.V. Klimov, S.V. Baranov, S.I. Allakhverdiev, Bicarbonate protects the donor side of
 photosystem II against photoinhibition and thermoinactivation, *FEBS Lett.* **418**
 (1997) 243–246.
- [51] K.N. Ferreira, T.M. Iverson, K. Maghlaoui, J. Barber, S. Iwata, Architecture of the
 photosynthetic oxygen-evolving center, *Science* **303** (2004) 1831–1838.
- [52] B. Loll, J. Kern, W. Saenger, A. Zouni, J. Biesiadka, Towards complete cofactor arrange-
 ment in the 3.0 Å resolution structure of photosystem II, *Nature* **438** (2005)
 1040–1044.
- [53] Govindjee, H.G. Weger, D.H. Turpin, J.J.S. van Rensen, O.J. Devos, J.F.H. Snel, Formate re-
 leases carbon dioxide/bicarbonate from thylakoid membranes – measurements by
 mass spectroscopy and infrared gas analyzer, *Naturwissenschaften* **78** (1991)
 168–170.
- [54] D. Shevela, V. Klimov, J. Messinger, Interactions of photosystem II with bicarbon-
 ate, formate and acetate, *Photosynth. Res.* **94** (2007) 247–264.
- [55] G. Ulas, G. Olack, G.W. Brudvig, Evidence against bicarbonate bound in the O₂-evolving
 complex of photosystem II, *Biochemistry* **47** (2008) 3073–3075.
- [56] P.E.M. Siegbahn, M. Lundberg, Hydroxide instead of bicarbonate in the structure
 of the oxygen evolving complex, *J. Inorg. Biochem.* **100** (2006) 1035–1040.
- [57] E.M. Sproviero, J.A. Gascon, J.P. McEvoy, G.W. Brudvig, V.S. Batista, QM/MM
 models of the O₂-evolving complex of photosystem II, *J. Chem. Theory Comput.*
2 (2006) 1119–1134.
- [58] S.I. Allakhverdiev, T. Tsuchiya, K. Watabe, A. Kojima, D.A. Los, T. Tomo, V.V. Klimov,
 M. Mimuro, Redox potentials of primary electron acceptor quinone molecule (Q_A)
 and coserved energetics of photosystem II in cyanobacteria with chlorophyll *a* and
 chlorophyll *d*, *Proc. Natl. Acad. Sci. U. S. A.* **108** (2011) 8054–8058.
- [59] S.V. Baranov, G.M. Ananyev, V.V. Klimov, G.C. Dismukes, Bicarbonate accelerates
 assembly of the inorganic core of the water-oxidizing complex in manganese
 depleted photosystem II: a proposed biogeochemical role for atmospheric car-
 bon dioxide in oxygenic photosynthesis, *Biochemistry* **39** (2000) 6060–6065.
- [60] G.M. Ananyev, L. Zaltsman, C. Vasko, G.C. Dismukes, The inorganic biochemistry of
 photosynthetic oxygen evolution/water oxidation, *Biochim. Biophys. Acta* **1503**
 (2001) 52–68.
- [61] J. Dasgupta, G.M. Ananyev, G.C. Dismukes, Photoassembly of the water-oxidizing
 complex in photosystem II, *Coord. Chem. Rev.* **252** (2008) 347–360.

- 1374 [62] Y.N. Kozlov, S.K. Zharmukhamedov, K.G. Tikhonov, J. Dasgupta, A.A. Kazakova, G.C. Dismukes, V.V. Klimov, Oxidation potentials and electron donation to photosystem II of manganese complexes containing bicarbonate and carboxylate ligands, *Phys. Chem. Chem. Phys.* 6 (2004) 4905–4911.
- 1375 [63] Y.N. Kozlov, K.G. Tikhonov, O.M. Zastrizhnaya, V.V. Klimov, pH dependence of the composition and stability of Mn^{III}-bicarbonate complexes and its implication for redox interaction of Mn^{II} with photosystem II, *J. Photochem. Photobiol. B Biol.* 101 (2010) 362–366.
- 1376 [64] J. Dasgupta, A.M. Tyrishkin, Y.N. Kozlov, V.V. Klimov, G.C. Dismukes, Carbonate complexation of Mn²⁺ in the aqueous phase: redox behavior and ligand binding modes by electrochemistry and EPR spectroscopy, *J. Phys. Chem. B* 110 (2006) 5099–5111.
- 1377 [65] J. Dasgupta, A.M. Tyrishkin, G.C. Dismukes, ESEEM spectroscopy reveals carbonate and an N-donor protein–ligand binding to Mn²⁺ in the photoassembly reaction of the Mn₄Ca cluster in photosystem II, *Angew. Chem. Int. Ed.* 46 (2007) 8028–8031.
- 1378 [66] J. Dasgupta, A.M. Tyrishkin, S.V. Baranov, G.C. Dismukes, Bicarbonate coordinates to Mn³⁺ during photo-assembly of the catalytic Mn₄Ca core of photosynthetic water oxidation: EPR characterization, *Appl. Magn. Reson.* 37 (2010) 137–150.
- 1379 [67] O.V. Pobeguts, T.N. Smolova, O.M. Zastrizhnaya, V.V. Klimov, Protective effect of bicarbonate against extraction of the extrinsic proteins of the water-oxidizing complex from photosystem II membrane fragments, *Biochim. Biophys. Acta* 1767 (2007) 624–632.
- 1380 [68] O.V. Pobeguts, T.N. Smolova, D.S. Timoshevsky, V.V. Klimov, Interaction of bicarbonate with the manganese-stabilizing protein of photosystem II, *J. Photochem. Photobiol. B Biol.* 100 (2010) 30–37.
- 1381 [69] T.M. Bricker, L.K. Frankel, Auxiliary functions of the PsbO, PsbP and PsbQ proteins of higher plant photosystem II: a critical analysis, *J. Photochem. Photobiol. B Biol.* 104 (2011) 165–178.
- 1382 [70] N. Nelson, C.F. Yocum, Structure and function of photosystems I and II, *Annu. Rev. Plant Biol.* 57 (2006) 521–565.
- 1383 [71] L.I. Kristsalik, Energetics of multielectron reactions. Photosynthetic oxygen evolution, *Biochim. Biophys. Acta* 849 (1986) 162–171.
- 1384 [72] O.V. Moskvina, T.V. Shutova, M.S. Khristin, L.K. Ignatova, A. Villarejo, G. Samuelsson, V.V. Klimov, B.N. Ivanov, Carbonic anhydrase activities in pea thylakoids – a photosystem II core complex-associated carbonic anhydrase, *Photosynth. Res.* 79 (2004) 93–100.
- 1385 [73] A.V. Shitov, O.V. Pobeguts, T.N. Smolova, S.I. Allakhverdiev, V.V. Klimov, Manganese-dependent carbonic anhydrase activity of photosystem II proteins, *Biochemistry (Mosc.)* 74 (2009) 509–517.
- 1386 [74] Y.K. Lu, A.J. Stemler, Extrinsic photosystem II carbonic anhydrase in maize mesophyll chloroplasts, *Plant Physiol.* 128 (2002) 643–649.
- 1387 [75] M.S. Khristin, L.K. Ignatova, N.N. Rudenko, B.N. Ivanov, V.V. Klimov, Photosystem II associated carbonic anhydrase activity in higher plants is situated in core complex, *FEBS Lett.* 577 (2004) 305–308.
- 1388 [76] Y.K. Lu, S.M. Theg, A.J. Stemler, Carbonic anhydrase activity of the photosystem II OEC33 protein from pea, *Plant Cell Physiol.* 46 (2005) 1944–1953.
- 1389 [77] I.L. McConnell, M.R. Badger, T. Wydrzynski, W. Hillier, A quantitative assessment of the carbonic anhydrase activity in photosystem II, *Biochim. Biophys. Acta* 1767 (2007) 639–647.
- 1390 [78] Y.K. Lu, A.J. Stemler, Differing responses of the two forms of photosystem II carbonic anhydrase to chloride, cations, and pH, *Biochim. Biophys. Acta* 1767 (2007) 633–638.
- 1391 [79] A. Stemler, Carbonic Anhydrase: Molecular Insights Applied to Photosystem II Research in Thylakoid Membranes, *The American Society of Plant Physiologists*, Washington, 1985.
- 1392 [80] M. Moubarak-Milad, A. Stemler, Oxidation-reduction potential dependence of photosystem II carbonic anhydrase in maize thylakoids, *Biochemistry* 33 (1994) 4432–4438.
- 1393 [81] A.J. Stemler, The case for chloroplast thylakoid carbonic anhydrase, *Physiol. Plant.* 99 (1997) 348–353.
- 1394 [82] A.V. Shitov, S.K. Zharmukhamedov, T. Shutova, S.I. Allakhverdiev, G. Samuelsson, V.V. Klimov, A carbonic anhydrase inhibitor induces bicarbonate-reversible suppression of electron transfer in pea photosystem 2 membrane fragments, *J. Photochem. Photobiol. B Biol.* 104 (2011) 366–371.
- 1395 [83] A.N. Tikhonov, R.V. Agafonov, I.A. Grigor'ev, I.A. Kirilyuk, V.V. Ptushenko, B.V. Trubitsin, Spin-probes designed for measuring the intrathylakoid pH in chloroplasts, *Biochim. Biophys. Acta* 1777 (2008) 285–294.
- 1396 [84] P.A. Joliot, G. Finazzi, Proton equilibration in the chloroplast modulates multiphasic kinetics of nonphotochemical quenching of fluorescence in plants, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 12728–12733.
- 1397 [85] D. Carrieri, G. Ananyev, T. Brown, G.C. Dismukes, *In vivo* bicarbonate requirement for water oxidation by photosystem II in the hypercarbonate-requiring cyanobacterium *Arthrospira maxima*, *J. Inorg. Biochem.* 101 (2007) 1865–1874.
- 1398 [86] G. Ananyev, G.C. Dismukes, How fast can photosystem II split water? Kinetic performance at high and low frequencies, *Photosynth. Res.* 84 (2005) 355–365.
- 1399 [87] F.P. Boyle, Some factors involved in oxygen evolution from tritiated spinach leaves, *Science* 108 (1948) 359–360.
- 1400 [88] O. Warburg, G. Krippahl, Notwendigkeit der Kohlensäure für die Chinon- und Ferricyanid-Reaktionen in Grünen Grana, *Z. Naturforsch. B* 15 (1960) 367–369.
- 1401 [89] F.B. Abeles, A.H. Brown, B.C. Mayne, Stimulation of the Hill reaction by carbon dioxide, *Plant Physiol.* 36 (1961) 202–207.
- 1402 [90] B.K. Stern, B. Vennesland, The carbon dioxide requirement for photoevolution of oxygen by chloroplast preparations, *J. Biol. Chem.* 235 (1960) 51–53.
- 1403 [91] B.K. Stern, B. Vennesland, The effect of carbon dioxide on the Hill reaction, *J. Biol. Chem.* 237 (1962) 596–602.
- 1404 [92] B. Vennesland, The stimulation of the Hill reaction by CO₂, in: B. Kok, A.T. Jagendorf (Eds.), *Photosynthetic Mechanisms of Green Plants*, National Academy of Science – National Research Council, Washington, DC, 1963, pp. 424–427.
- 1405 [93] S. Izawa, Stimulatory effects of carbon dioxide upon the Hill reaction as observed with the addition of carbonic anhydrase to reaction mixture, *Plant Cell Physiol.* 3 (1962) 221–227.
- 1406 [94] J.J. Heise, H. Gaffron, Catalytic effects of carbon dioxide in carbon dioxide assimilating cells, *Plant Cell Physiol.* 4 (1963) 1–11.
- 1407 [95] N.E. Good, Carbon dioxide and the Hill reaction, *Plant Physiol.* 38 (1963) 298–304.
- 1408 [96] N.E. Good, Interpretations of the carbon dioxide dependence of the Hill reaction, *Can. J. Bot.* 43 (1965) 119–127.
- 1409 [97] T. Punnett, R.V. Iyer, The enhancement of photophosphorylation and the Hill reaction by carbon dioxide, *J. Biol. Chem.* 239 (1964) 2335–2339.
- 1410 [98] T. Punnett, Influence of growth conditions on the enhancement of photophosphorylation by carbon dioxide, *Plant Physiol.* 40 (1965) 1283–1284.
- 1411 [99] P.P. Batra, A.T. Jagendorf, Bicarbonate effects on the Hill reaction and photophosphorylation, *Plant Physiol.* 40 (1965) 1074–1079.
- 1412 [100] J. West, R. Hill, Carbon dioxide and the reduction of indophenol and ferricyanide by chloroplasts, *Plant Physiol.* 42 (1967) 819–826.
- 1413 [101] A. Stemler, Govindjee, Bicarbonate stimulation of oxygen evolution, ferricyanide reduction and photoinactivation using isolated chloroplasts, *Plant Cell Physiol.* 15 (1974) 533–544.
- 1414 [102] A. Stemler, Govindjee, Effects of bicarbonate ion on chlorophyll *a* fluorescence transients and delayed light-emission from maize chloroplasts, *Photochem. Photobiol.* 19 (1974) 227–232.
- 1415 [103] P. Jursinic, J. Warden, Govindjee, A major site of bicarbonate effect in system II reaction. Evidence from ESR signal II *vf*, fast fluorescence yield changes and delayed light emission, *Biochim. Biophys. Acta* 440 (1976) 322–330.
- 1416 [104] Govindjee, M.P.J. Pulles, R. Govindjee, H.J. van Gorkom, L.N.M. Duysens, Inhibition of reoxidation of secondary-electron acceptor of photosystem II by bicarbonate depletion, *Biochim. Biophys. Acta* 449 (1976) 602–605.
- 1417 [105] R. Khanna, Govindjee, T. Wydrzynski, Site of bicarbonate effect in Hill reaction. Evidence from the use of artificial electron acceptors and donors, *Biochim. Biophys. Acta* 462 (1977) 208–214.
- 1418 [106] U. Siggel, R. Khanna, G. Renger, Govindjee, Investigation of absorption changes of plastoquinone system in broken chloroplasts: effect of bicarbonate depletion, *Biochim. Biophys. Acta* 462 (1977) 196–207.
- 1419 [107] R. Khanna, R. Wagner, W. Junge, Govindjee, Effects of CO₂-depletion on proton uptake and release in thylakoid membranes, *FEBS Lett.* 121 (1980) 222–224.
- 1420 [108] R. Khanna, K. Pfister, A. Keresztes, J.J. van Rensen, Govindjee, Evidence for a close spatial location of the binding sites for CO₂ and for photosystem II inhibitors, *Biochim. Biophys. Acta* 634 (1981) 105–116.
- 1421 [109] W.F.J. Vermaas, J.J. Van Rensen, Govindjee, The interaction between bicarbonate and the herbicide ioxynil in the thylakoid membrane and the effects of amino acid modification on bicarbonate action, *Biochim. Biophys. Acta* 681 (1982) 242–247.
- 1422 [110] J.J. Eaton-Rye, Govindjee, A study of the specific effect of bicarbonate on photosynthetic electron transport in the presence of methyl viologen, *Photobiophys.* 8 (1984) 279–288.
- 1423 [111] H.H. Robinson, J.J. Eaton-Rye, J.J.S. Van Rensen, Govindjee, The effects of bicarbonate depletion and formate incubation on the kinetics of oxidation–reduction reactions of the photosystem II quinone acceptor complex, *Z. Naturforsch. C* 39 (1984) 382–385.
- 1424 [112] Govindjee, H.Y. Nakatani, A.W. Rutherford, Y. Inoue, Evidence from thermoluminescence for bicarbonate action on the recombination reactions involving the secondary quinone electron acceptor of photosystem II, *Biochim. Biophys. Acta* 766 (1984) 416–423.
- 1425 [113] R.E. Zeebe, D.A. Wolf-Gladrow, CO₂ in seawater: equilibrium, kinetics, isotopes, in: D. Halpern (Ed.), *Elsevier Oceanography Series*, Elsevier, Amsterdam, 2001, p. 346.
- 1426 [114] D.J. Blubaugh, Govindjee, Bicarbonate, not CO₂, is the species required for the stimulation of photosystem II electron transport, *Biochim. Biophys. Acta* 848 (1986) 147–151.
- 1427 [115] G. Sarojini, Govindjee, On the active species in bicarbonate stimulation of Hill reaction in thylakoid membranes, *Biochim. Biophys. Acta* 634 (1981) 340–343.
- 1428 [116] D.J. Blubaugh, Govindjee, Kinetics of the bicarbonate effect and the number of bicarbonate-binding sites in thylakoid membranes, *Biochim. Biophys. Acta* 936 (1988) 208–214.
- 1429 [117] J.J. Eaton-Rye, Govindjee, Electron transfer through the quinone acceptor complex of photosystem II in bicarbonate-depleted spinach thylakoid membranes as a function of actinic flash number and frequency, *Biochim. Biophys. Acta* 935 (1988) 237–247.
- 1430 [118] J.J. Eaton-Rye, Govindjee, Electron transfer through the quinone acceptor complex of photosystem II after one or two actinic flashes in bicarbonate-depleted spinach thylakoid membranes, *Biochim. Biophys. Acta* 935 (1988) 248–257.
- 1431 [119] J. Cao, Govindjee, Bicarbonate effect on electron flow in cyanobacterium *Synechocystis* PCC 6803, *Photosynth. Res.* 19 (1988) 277–285.
- 1432 [120] G. Garab, Z. Rozsa, Govindjee, Carbon dioxide affects charge accumulation in leaves: measurements by thermoluminescence, *Naturwissenschaften* 75 (1988) 517–519.
- 1433 [121] Govindjee, H. Robinson, A.R. Crofts, J.J.S. Van Rensen, Bicarbonate does not influence electron transfer to the reaction center chlorophyll *a* of photosystem II: measurements by chlorophyll *a* fluorescence rise in microseconds, *Naturwissenschaften* 76 (1989) 119–121.
- 1434 [122] R.J. Shopes, D.J. Blubaugh, C.A. Wraight, Govindjee, Absence of a bicarbonate-depletion effect in electron transfer between quinones in chromatophores and reaction centers of *Rhodospirillum rubrum*, *Biochim. Biophys. Acta* 974 (1989) 114–118.
- 1435 [123] X. Wang, J. Cao, P. Maroti, H.U. Stiltz, U. Finkele, C. Lauterwasser, W. Zinth, D. Oesterhelt, Govindjee, C.A. Wraight, Is bicarbonate in photosystem II the

- equivalent of the glutamate ligand to the iron atom in bacterial reaction centers? *Biochim. Biophys. Acta* 1100 (1992) 1–8.
- [124] H. Michel, J. Deisenhofer, Relevance of the photosynthetic reaction center from purple bacteria to the structure of photosystem II, *Biochemistry* 27 (1988) 1–7.
- [125] Govindjee, C. Vernotte, B. Peteri, C. Astier, A.L. Etienne, Differential sensitivity of bicarbonate-reversible formate effects on herbicide-resistant mutants of *Synechocystis* 6714, *FEBS Lett.* 267 (1990) 273–276.
- [126] A. Stemler, Absence of a formate-induced release of bicarbonate from photosystem 2, *Plant Physiol.* 91 (1989) 287–290.
- [127] J. Oscar, O.J. De Vos, J.J.S. Van Rensen, Govindjee, Photosystem II electron flow requires bound bicarbonate, in: P. Mathis (Ed.), *Photosynthesis: From Light to Biosphere*, Kluwer Academic Publishers, The Netherlands, 1995, pp. 567–570.
- [128] P.A. Jursinic, A. Stemler, High-rates of photosystem II electron flow occur in maize thylakoids when the high-affinity binding-site for bicarbonate is empty of all monovalent anions or has bicarbonate bound, *Biochim. Biophys. Acta* 1098 (1992) 359–367.
- [129] Govindjee, B. Schwarz, J.D. Rochaix, R.J. Strasser, The herbicide-resistant D1 mutant L275F of *Chlamydomonas reinhardtii* fails to show the bicarbonate-reversible formate effect on chlorophyll *a* fluorescence transients, *Photosynth. Res.* 27 (1991) 199–208.
- [130] C.H. Xu, S. Taoka, A.R. Crofts, Govindjee, Kinetic characteristics of formate formic-acid binding at the plastoquinone reductase site in spinach thylakoids, *Biochim. Biophys. Acta* 1098 (1991) 32–40.
- [131] J.C. Cao, W.F.J. Vermaas, Govindjee, Arginine residues in the D2 polypeptide may stabilize bicarbonate binding in photosystem II of *Synechocystis* sp PCC 6803, *Biochim. Biophys. Acta* 1059 (1991) 171–180.
- [132] Govindjee, P. Eggenberg, K. Pfister, R.J. Strasser, Chlorophyll *a* fluorescence yield decay in herbicide-resistant D1 mutant of *Chlamydomonas reinhardtii* and the formate effect, *Biochim. Biophys. Acta* 1101 (1992) 353–358.
- [133] R.J. Strasser, P. Eggenberg, K. Pfister, Govindjee, An equilibrium model for electron transfer in photosystem II acceptor complex: an application to *Chlamydomonas reinhardtii* cells of D1 mutants and those treated with formate, *Arch. Science Geneve* 45 (1992) 207–224.
- [134] J. Cao, N. Ohad, J. Hirschberg, J. Xiong, Govindjee, Binding affinity of bicarbonate and formate in herbicide-resistant D1 mutants of *Synechococcus* sp. PCC 7942, *Photosynth. Res.* 34 (1992) 397–408.
- [135] Govindjee, Bicarbonate-reversible inhibition of plastoquinone reductase in photosystem II, *Z. Naturforsch.* C 48 (1993) 251–258.
- [136] P. Mäenpää, T. Miranda, E. Tyystjärvi, T. Tyystjärvi, Govindjee, J.M. Ducruet, A.L. Etienne, D. Kirilovsky, A mutation in the D-de loop of D-1 modifies the stability of the $S_2Q_A^-$ and $S_2Q_B^-$ states in photosystem II, *Plant Physiol.* 107 (1995) 187–197.
- [137] R.S. Hutchison, J. Xiong, R.T. Sayre, Govindjee, Construction and characterization of a photosystem II D1 mutant (arginine-269-glycine) of *Chlamydomonas reinhardtii*, *Biochim. Biophys. Acta* 1277 (1996) 83–92.
- [138] J. Xiong, S. Subramaniam, Govindjee, Modeling of the D1/D2 proteins and cofactors of the photosystem II reaction center: implications for herbicide and bicarbonate binding, *Protein Sci.* 5 (1996) 2054–2073.
- [139] S. Rose, J. Minagawa, M. Seufferheld, S. Padden, B. Svensson, D.R. Kolling, A.R. Crofts, Govindjee, D1-arginine257 mutants (R257E, K, and Q) of *Chlamydomonas reinhardtii* have a lowered Q_B redox potential: analysis of thermoluminescence and fluorescence measurements, *Photosynth. Res.* 98 (2008) 449–468.
- [140] J. Xiong, J. Minagawa, A. Crofts, Govindjee, Loss of inhibition by formate in newly constructed photosystem II D1 mutants, D1-R257E and D1-R257M, of *Chlamydomonas reinhardtii*, *Biochim. Biophys. Acta* 1365 (1998) 473–491.
- [141] V. Petrouleas, A.R. Crofts, The iron-quinone acceptor complex, in: T. Wydrzynski, K. Satoh (Eds.), *Photosystem II. The Light-Driven Water:Plastoquinone Oxidoreductase*, Springer, Dordrecht, 2005, pp. 177–206.
- [142] S. Padden, J. Minagawa, A. Kanazawa, Govindjee, A.R. Crofts, Site specific mutagenesis reveals a critical role for histidine 252 of the D1 subunit in the two-electron gate of photosystem II, 28th Annual Eastern Regional Photosynthesis Conference, Marine Biological Laboratory, Woods Hole MA, USA, 2011.
- [143] W.F.J. Vermaas, A.W. Rutherford, EPR measurements of the effects of bicarbonate and triazine resistance on the acceptor side of photosystem II, *FEBS Lett.* 175 (1984) 243–248.
- [144] B.A. Diner, V. Petrouleas, Light-induced oxidation of the acceptor-side Fe(II) of photosystem II by exogenous quinones acting through the Q_B binding-site. 2. Blockage by inhibitors and their effects on the Fe(III) EPR spectra, *Biochim. Biophys. Acta* 893 (1987) 138–148.
- [145] J.H.A. Nugent, A.R. Corrie, C. Demetriou, M.C.W. Evans, C.J. Lockett, Bicarbonate binding and the properties of photosystem II electron acceptors, *FEBS Lett.* 235 (1988) 71–75.
- [146] V. Petrouleas, B.A. Diner, Identification of Q_{400} , a high potential electron acceptor of photosystem II, with the iron of the quinone-iron acceptor complex, *Biochim. Biophys. Acta* 849 (1986) 264–275.
- [147] W.F.J. Vermaas, Govindjee, Bicarbonate or CO_2 as a requirement for efficient electron transport on the acceptor side of photosystem II, in: Govindjee (Ed.), *Photosynthesis. Development, Carbon Metabolism, and Plant Productivity*, Academic Press, New York, 1982, pp. 541–558.
- [148] B.A. Diner, V. Petrouleas, J.J. Wendoloski, The iron-quinone electron acceptor complex of photosystem II, *Physiol. Plant.* 81 (1991) 423–436.
- [149] R. Hienerwadel, C. Berthomieu, Bicarbonate binding to the non-heme iron of photosystem II investigated by Fourier transform infrared difference spectroscopy and ^{13}C -labeled bicarbonate, *Biochemistry* 34 (1995) 16288–16297.
- [150] C. Berthomieu, R. Hienerwadel, Iron coordination in photosystem II: interaction between bicarbonate and Q_B pocket studied by Fourier transform infrared spectroscopy, *Biochemistry* 40 (2001) 4044–4052.
- [151] D. Shevela, V. Klimov, J. Messinger, Formate-induced release of carbon dioxide/hydrogencarbonate from photosystem II, in: J.F. Allen, E. Gannt, J.H. Golbeck, B. Osmond (Eds.), *Photosynthesis. Energy from the Sun: 14th International Congress on Photosynthesis*, Springer, Glasgow, 2008, pp. 497–501.
- [152] Y.M. Feyziev, D. Yoneda, T. Yoshii, N. Katsuta, A. Kawamori, Y. Watanabe, Formate-induced inhibition of the water-oxidizing complex of photosystem II studied by EPR, *Biochemistry* 39 (2000) 3848–3855.
- [153] N. Cox, L. Jin, A. Jaszewski, P.J. Smith, E. Krausz, A.W. Rutherford, R. Pace, The semiquinone-iron complex of photosystem II: structural insights from ESR and theoretical simulation; evidence that the native ligand to the non-heme iron is carbonate, *Biophys. J.* 97 (2009) 2024–2033.
- [154] R. Takahashi, A. Boussac, M. Sugiura, T. Noguchi, Structural coupling of a tyrosine side chain with the non-heme iron center in photosystem II as revealed by light-induced Fourier transform infrared difference spectroscopy, *Biochemistry* 48 (2009) 8994–9001.
- [155] A. Sedoud, L. Kastner, N. Cox, S. El-Alaoui, D. Kirilovsky, A.W. Rutherford, Effects of formate binding on the quinone-iron electron acceptor complex of photosystem II, *Biochim. Biophys. Acta* 1807 (2011) 216–226.
- [156] P. Chernev, I. Zaharieva, H. Dau, M. Haumann, Carboxylate shifts steer inter-quinone electron transfer in photosynthesis, *J. Biol. Chem.* 286 (2011) 5368–5374.
- [157] S. Hermes, O. Bremm, F. Garczarek, V. Derrien, P. Liebisch, P. Loja, P. Sebban, K. Gerwert, M. Haumann, A time-resolved iron-specific X-ray absorption experiment yields no evidence for an $Fe^{2+} \rightarrow Fe^{3+}$ transition during $Q_A \rightarrow Q_B$ electron transfer in the photosynthetic reaction center, *Biochemistry* 45 (2006) 353–359.
- [158] F. Müh, C. Glöckner, J. Hellmich, A. Zouni, Light-induced quinone reduction in photosystem II, *Biochim. Biophys. Acta* 1817 (2012) 44–65.
- [159] Govindjee, J.-R. Shen, A.R. Crofts, Photosystem II: unique role of bicarbonate, International Conference "Photosynthesis Research for Sustainability", Baku, Azerbaijan, 2011.
- [160] A. Zouni, H.T. Witt, J. Kern, P. Fromme, N. Krauß, W. Saenger, P. Orth, Crystal structure of photosystem II from *Synechococcus elongatus* at 3.8 Å resolution, *Nature* 409 (2001) 739–743.
- [161] N. Kamiya, J.-R. Shen, Crystal structure of oxygen-evolving photosystem II from *Thermosynechococcus vulcanus* at 3.7 Å resolution, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 98–103.
- [162] A. Guskov, J. Kern, A. Gabdulkhakov, M. Broser, A. Zouni, W. Saenger, Cyanobacterial photosystem II at 2.9-Ångstrom resolution and the role of quinones, lipids, channels and chloride, *Nat. Struct. Mol. Biol.* 16 (2009) 334–342.
- [163] J. Yano, J. Kern, K.D. Irrgang, M.J. Latimer, U. Bergmann, P. Glatzel, Y. Pushkar, J. Biesiadka, B. Loll, K. Sauer, J. Messinger, A. Zouni, V.K. Yachandra, X-ray damage to the Mn_4Ca complex in single crystals of photosystem II: a case study for metalloprotein crystallography, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 12047–12052.
- [164] H. Fukuzawa, T. Ogawa, A. Kaplan, The uptake of CO_2 by cyanobacteria and microalgae, in: J.J. Eaton-Rye, B.C. Tripathy, T.D. Sharkey (Eds.), *Photosynthesis: Plastid Biology, Energy Conversion and Respiration*, Springer, Dordrecht, 2012, pp. 625–650.
- [165] P.G. Falkowski, Tracing oxygen's imprint on Earth's metabolic evolution, *Science* 311 (2006) 1724–1725.
- [166] L.P. Kump, The rise of atmospheric oxygen, *Nature* 451 (2008) 277–278.
- [167] M.F. Hohmann-Marriott, R.E. Blankenship, Evolution of photosynthesis, *Annu. Rev. Plant Biol.* 62 (2011) 515–548.
- [168] R.E. Blankenship, Early evolution of photosynthesis, *Plant Physiol.* 154 (2010) 434–438.
- [169] Govindjee, D. Shevela, Adventures with cyanobacteria: a personal perspective, *Front. Plant Sci.* 2 (2011) 28, <http://dx.doi.org/10.3389/fpls.2011.00028>.
- [170] G.C. Dismukes, V.V. Klimov, S.V. Baranov, Y.N. Kozlov, J. DasGupta, A. Tyrtyshkin, The origin of atmospheric oxygen on earth: the innovation of oxygenic photosynthesis, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 2170–2175.
- [171] A.A. Khorobrykh, V.V. Terentyev, S.K. Zharmukhamedov, V.V. Klimov, Redox interaction of Mn-bicarbonate complexes with reaction centres of purple bacteria, *Philos. Trans. R. Soc. Lond. B* 363 (2008) 1245–1251.