

Mayumi Tsuchimoto · Shinobu Yasuo ·
Masahiro Funada · Makoto Aoki · Hiromi Sasagawa ·
Takashi Yoshimura · Osamu Tadauchi ·
Sydney A. Cameron · Yasuo Kitagawa ·
Tatsuhiko Kadowaki

Conservation of novel *Mahya* genes shows the existence of neural functions common between Hymenoptera and Deuterostome

Received: 28 May 2005 / Accepted: 9 August 2005
© Springer-Verlag 2005

Abstract Honeybees have been shown to exhibit cognitive performances that were thought to be specific to some vertebrates. However, the molecular and cellular mechanisms of such cognitive abilities of the bees have not been understood. We have identified a novel gene, *Mahya*, expressed in the brain of the honeybee, *Apis mellifera*, and

other Hymenoptera. *Mahya* orthologues are present in Deuterostomes but are absent or highly diverged in nematodes and, intriguingly, in two dipteran insects (fruit fly and mosquito) and Lepidoptera (silk moth). *Mahya* genes encode novel secretory proteins with a follistatin-like domain (Kazal-type serine/threonine protease inhibitor domain and EF-hand calcium-binding domain), two immunoglobulin domains, and a C-terminal novel domain. Honeybee *Mahya* is expressed in the mushroom bodies and antennal lobes of the brain. Zebra fish *Mahya* orthologues are expressed in the olfactory bulb, telencephalon, habenula, optic tectum, and cerebellum of the brain. Mouse *Mahya* orthologues are expressed in the olfactory bulb, hippocampus, and cerebellum of the brain. These results suggest that *Mahya* may be involved in learning and memory and in processing of sensory information in Hymenoptera and vertebrates. Furthermore, the limited existence of *Mahya* in the genomes of Hymenoptera and Deuterostomes supports the hypothesis that the genes typically represented by *Mahya* were lost or highly diverged during the evolution of the central nervous system of specific Bilaterian branches under the specific selection and subsequent adaptation associated with different ecologies and life histories.

Communicated by C. Desplan

M. Tsuchimoto · M. Funada · M. Aoki · Y. Kitagawa ·
T. Kadowaki (✉)
Department of Bioengineering Sciences,
Graduate School of Bioagricultural Sciences,
Nagoya University,
Chikusa,
Nagoya, 464-8601, Japan
e-mail: emi@nuagr1.agr.nagoya-u.ac.jp
Tel.: +81-52-7895237
Fax: +81-52-7895237

S. Yasuo · T. Yoshimura
Division of Biomodeling,
Graduate School of Bioagricultural Sciences,
Nagoya University,
Chikusa,
Nagoya, 464-8601, Japan

Present address:

M. Aoki
Laboratory for Developmental Gene Regulation,
Brain Science Institute,
The Institute of Physical and Chemical Research (RIKEN),
Saitama, 351-0198, Japan

H. Sasagawa
Foundation for Advancement for International Science,
Tsukuba, 305-0062, Japan

O. Tadauchi
Graduate School of Bioresource and Bioenvironmental
Sciences, Kyushu University,
Higashi-ku,
Fukuoka, 812-8581, Japan

S. A. Cameron
Department of Entomology and Program in Ecology and
Evolutionary Biology, University of Illinois,
Urbana, IL 61801, USA

Keywords Comparative genomics · Evolution ·
Gene loss · CNS · Hymenoptera

Introduction

The honeybee, *Apis mellifera*, belongs to Hymenoptera, one of four large holometabolous insect orders. Honeybees are highly social insects and have been used as a model system to study complex animal behavior in addition to learning and memory. Honeybees form colonies consisting of a single queen, hundreds of males, and thousands of female worker bees. In contrast to the queen and male bees, workers perform a wide variety of tasks to maintain the colony, including nursing and foraging for food. In

comparison to performing tasks inside the dark hive, foraging involves extensive flight, learning and memorizing food locations, navigation using multiple cues (e.g., sun compass), and communication by a dance language (Fahrbach and Robinson 1995; Menzel and Muller 1996; Menzel and Giurfa 2001). These tasks may require better modules for sensory-input processing and enhanced learning and memory capabilities than other insects. The social behavior and cognitive ability of honeybees partially match those of higher vertebrates, a recognition that led to the launching of the honeybee genome project.

Cross-species comparison of genome sequences and EST data sets has given us enormous insight into the evolution of metazoan genomes. Regarding the origin and evolution of the central nervous system (CNS) of Protostomes and Deuterostomes, most of the nervous system-related genes of the planarian *Dugesia japonica* (Platyhelminthes, Tricladida) are shared with fruit fly, nematode, and human. This suggests at least the presence of a common ancestral CNS at the molecular level (Mineta et al. 2003). However, a few planarian genes are not conserved among the above animals, suggesting that gene loss occurred during the evolution of the Bilaterian CNS. This is also consistent with the high evolutionary rates of model invertebrates, fruit fly, and nematode compared to those of vertebrates (Mushegian et al. 1998; Zdobnov et al. 2002). Research on *Acropora millepora* (Anthozoan, Cnidaria) supports a view that a significant proportion of the genes present only in the vertebrates and absent in the model invertebrates is not a vertebrate-specific invention. Instead, these genes appear to have been lost in the specific metazoan lineages that branched from the common ancestors during evolution (Kortschak et al. 2003). The comparative analysis of honeybee EST data set with genome databases also demonstrates that over 100 honeybee cDNA sequences conserved with other organisms are absent in the fruit fly genome (Whitfield et al. 2002). Meanwhile, a number of studies examining genes involved in morphogenesis have revealed the spectacular conservation of the developmental programs between fruit fly and the vertebrates. These results thus demonstrate that the genes functioning for basic biological programs (e.g., the process of embryogenesis) were conserved across the metazoan species, but some genes were either generated or lost in the particular metazoan branches under positive and negative selection pressures.

In addition to two dipteran insects, *Drosophila melanogaster* and *Anopheles gambiae*, the genome projects of *Apis mellifera*, *Bombyx mori* (silk moth, Lepidoptera), and *Tribolium castaneum* (red flour beetle, Coleoptera) promise new insights into how the genomes of holometabolous insects evolved in comparison to other metazoans. While most of their genes are expected to be conserved, there will be some genes that are not shared among these five model insect species but are present in other metazoans. Characterization of these genes (their distribution across Bilateria and Cnidaria and their expression patterns) will give us a clearer picture of the ancient evolutionary states and the relationship between gene loss and evolutionary change.

Here, we report a novel gene, *Mahya*, that is conserved in Hymenoptera and Deuterostomes but is absent in fruit fly, mosquito, silk moth, and nematode. *Mahya* genes encode novel secretory proteins and are highly expressed in the specific brain regions of honeybee, zebra fish, and mouse where learning and memory and processing of sensory information take place. Thus, *Mahya* appears to be one of the CNS-related genes retained and evolved in specific Bilaterian branches during evolution.

Materials and methods

Identification and isolation of *Mahya* orthologues from honeybee, ascidian, zebra fish, mouse, and human

We first identified the honeybee *Mahya* during a screen of genes differentially expressed between newly emerged bee and the forager brains (Tsuchimoto et al. 2004). The partial *Mahya* cDNA clone was used as a probe to screen the honeybee brain-specific cDNA library supplied by G. E. Robinson (University of Illinois). The 2.8 kb cDNA containing 3' UTR was obtained, and the cDNA with 5' end sequence was then isolated by a 5' rapid amplification of cDNA end (RACE) method. The full-length *Mahya* cDNA (3,585 bp) was isolated by a reverse transcription (RT)-polymerase chain reaction (PCR) method with two primers (5' CGCGGGGAGTACACCCCAAGCCATCAC 3' and 5' GGCCACCGATAGGGTAACGATTAGG 3') and then sequenced (accession number: AB231585). To identify and clone zebra fish *Mahya* orthologues (*drMahya-1* and *drMahya-2*), the zebra fish genome and EST database (http://www.sanger.ac.uk/Project3/D_erio/) were searched by BLAST with the honeybee *Mahya* protein sequence as a query. We identified two different *Mahya* orthologues with the significant *E* values represented by one EST clone (accession number: BG739057) and several genome sequences. The primers for 5' and 3' RACE were then designed, and the cDNAs with 5' and 3' ends were isolated. The partial cDNAs filling the gaps between 5' and 3' RACE products were isolated by RT-PCR. All of these cDNAs were sequenced (accession number: AB231586 and AB231587). Human (accession numbers: NP064501 and AX135099) and mouse (accession numbers: NP848788 and NP796033) *Mahya* orthologues were identified through a nr database search by BLAST. The cDNA sequence of *mMahya-2* (NP796033) contains one base insertion in the ORF that is corrected by assembling the two different reads of the same cDNA. The ascidian *Mahya* orthologue was identified by searching a *Ciona intestinalis* genome database (<http://www.genome.jgi-psf.org/ciona4/ciona4.home.html>) by BLAST with the *mMahya-1* protein sequence as a query. The initial attempt was unsuccessful with the honeybee *Mahya* protein sequence as a query. Search for *Mahya* orthologues in other animal species was carried out by both BLASTP and TBLASTN analysis (with low complexity filter, expected value: 10) of their genome

sequences using honeybee, ascidian, zebra fish, and mouse Mahya protein sequences as queries.

Analysis of the expression pattern of honeybee *Mahya* by Northern blot and RT-PCR

PolyA⁺RNA was isolated from the brains of 20 newly emerged bees, 7-day-old in-hive bees, and 28-day-old foragers. Two micrograms of polyA⁺RNA was electrophoresed with 1% agarose gel containing 2.2 M formaldehyde and then transferred to a nylon membrane. After the fixation of RNA, membranes were hybridized with ³²P-labeled *Mahya* and *EF-1alpha* cDNAs in a buffer containing 0.5 M sodium phosphate (pH 7.2), 7% (w/v) SDS, and 1 mM EDTA (pH 7.0) at 68°C. The signal was detected and analyzed with a Bioimage analyzer BAS2000 (Fuji). The expression of *Mahya* in the brain, thorax, and abdomen of honeybee workers, and in the brains of males and queens, was analyzed by RT-PCR with polyA⁺RNA isolated from each sample and the following two primers: 5' GTAATACCAACAGATAAAAATCCTG 3' and 5' AC AAATTGGATTTGAAGAATATGAAA 3'. The expression of *beta actin* was also analyzed as a control with the following two primers: 5' AATGGCAACTGCTGCATCA TCCTCAAGCTT 3' and 5' GAGATCCACATCTGTTG GAAGGTGGACA 3'. The amplified DNA fragments were sequenced to confirm their identity.

In situ hybridization of the brain sections of honeybee, zebra fish, and mouse

In situ hybridization of honeybee (foragers) brain sections with digoxigenin-labeled riboprobes was carried out as described in Funada et al. (2004). For the preparation of riboprobes, the partial honeybee *Mahya* cDNA (57-3580) with T7 and SP6 promoters was first PCR amplified, and then the sense and antisense riboprobes were made with either T7 or SP6 RNA polymerase and rNTPs containing digoxigenin-UTP. Hybridization was carried out overnight at 62°C. High-stringency posthybridization washes were performed at the same temperature. The brain sections were blocked and then incubated with 500-fold diluted alkaline phosphatase (AP)-conjugated antidigoxigenin antibody overnight at 4°C. The brain sections were washed, and then the AP substrate solution containing nitroblue tetrazolium and 5-bromo-4-chlor-3-indoly-phosphate was added. The color development was monitored under a microscope and carried out for the same periods on the samples hybridized with the sense and antisense riboprobes. In situ hybridization of zebra fish and mouse brain sections was carried out as described in Yoshimura et al. (2000). The antisense and sense 45 mer oligonucleotides were made against the following sequences—*drMahya-1*: 5' GGTGTGTGGATCTGACGGACGCTTCCACCAGAA CCACTGCGAGCT 3'; *drMahya-2*: 5' CCTGTGTGCGG ATCTGACGGGAAACTCTACCAGACCACTGTGAGC




Fig. 1 Domain structure of Mahya and the phylogenetic relationship among Mahya proteins from different species. The schematic representation of Mahya is shown in **a**. Mahya contains the N-terminal signal sequence (*solid oval*), Kazal-type serine/threonine protease inhibitor domain (*Kazal*), EF-hand calcium-binding domain (*EFh*), two immunoglobulin domains found in cell adhesion molecule (*IgCAM*), and C-terminal novel domain (*Mahya domain*). Mahya proteins from all species examined share the same domain

structure. The phylogenetic relationship among Mahya orthologues is shown in **b**. The vertebrate Mahya-1 and Mahya-2 proteins cluster into the separate clades. Honeybee and ascidian Mahya proteins are diverged from each other and from the vertebrate Mahya proteins. The tree was constructed by the neighbor-joining method. The *scale bar* indicates percent divergence or distance between the sequences. *Branch lengths* indicate phylogenetic divergence. *Numbers at the nodes* of the tree are bootstrap values

Table 1 Identity and similarity of full length *Mahya* proteins from different species

Identity (%)	Similarity (%)							
	<i>Mahya</i>	<i>drMahya-1</i>	<i>drMahya-2</i>	<i>mMahya-1</i>	<i>mMahya-2</i>	<i>hMahya-1</i>	<i>hMahya-2</i>	<i>ciMahya</i>
<i>Mahya</i>		50	49	50	48	51	49	45
<i>drMahya-1</i>	32		69	81	72	82	73	50
<i>drMahya-2</i>	30	51		72	71	72	71	50
<i>mMahya-1</i>	32	69	53		73	94	74	52
<i>mMahya-2</i>	31	55	53	56		73	87	48
<i>hMahya-1</i>	33	69	52	91	56		74	52
<i>hMahya-2</i>	31	56	53	57	81	57		46
<i>ciMahya</i>	26	31	30	32	30	32	28	

3'; *mMahya-1*: 5' GCGGAGCTGGTGCTGTCCACAGA GTGTCGGTGCTGAACCCCGGGT 3'; and *mMahya-2*: 5' AGCTTGGACTATCTCTAGCACGCCAGGGGACC AGCCGACAGCCGG 3'. They were labeled with [³³P] dATP and terminal deoxyribonucleotidyl transferase. The sagittal sections (20- μ m thickness) of the brain were prepared with a Cryostat. Hybridization was carried out overnight at 42°C. Two high-stringency posthybridization washes were performed at 55°C. The sections were air-dried and exposed to a Biomax-MR film for 2 weeks. Following the exposure, each slide was dipped into type NTB2 autoradiography emulsion and developed 4 weeks later at 4°C. The sections were then observed with a microscope for the signal detection.

Analysis of *Mahya* in other bee species by genomic PCR

Mahya was originally identified in *Apis mellifera*, the only animal containing *Mahya* in the Protostomes. To obtain insight into a possible relationship between *Mahya* and a specific animal behavior, we further searched for *Mahya* in the genomes of various bee species that exhibit not only different morphology but also express a wide range of social behavior. Based on honeybee genome sequence, honeybee *Mahya* consists of 18 exons, and exons 11–17 encode a novel C-terminal half domain (Mahya domain). Five different sets of primers (28–30 mer length; T_m at 49–

53.6°C) corresponding to the 5' and 3' end sequences of each of exons 13–17 were used for genomic PCR. These exon sequences are absent in the genomes of fruit fly, mosquito, and nematode. Genomic DNA was isolated from bee samples, either fresh, frozen, or stored in absolute ethanol, by the methods as described in Ashburner (1989). PCR was carried out (annealing temperature at 45°C; number of cycles, 40), and PCR products were analyzed by 2% agarose gel electrophoresis. Any lack of amplification could be due to (1) large mismatches between the primer and genome sequences, (2) different exon–intron organization of *Mahya* between honeybee and the other bees (the insertion of large intron sequences in the particular exons), or (3) absence of particular exons in the genomes. The PCR-amplified bands of expected sizes were extracted from the gel and sequenced to confirm their identity as exons encoding the Mahya domain. The sequences were identical to those of *Apis mellifera*, except for base substitutions at several positions.

Results

Identification of *Mahya* from honeybee, ascidian, zebra fish, mouse, and human

The longest open-reading frame of the honeybee *Mahya* cDNA encodes a 101-kDa protein of 898 amino acids with

Table 2 Identity and similarity of the Mahya domains of *Mahya* proteins from different species

Identity (%)	Similarity (%)							
	<i>Mahya</i>	<i>drMahya-1</i>	<i>drMahya-2</i>	<i>mMahya-1</i>	<i>mMahya-2</i>	<i>hMahya-1</i>	<i>hMahya-2</i>	<i>ciMahya</i>
<i>Mahya</i>		55	56	58	53	57	53	45
<i>drMahya-1</i>	33		70	84	72	85	72	52
<i>drMahya-2</i>	32	48		71	72	71	70	50
<i>mMahya-1</i>	34	71	47		73	95	72	51
<i>mMahya-2</i>	32	53	49	54		74	90	45
<i>hMahya-1</i>	34	72	46	91	54		74	51
<i>hMahya-2</i>	31	53	48	54	83	56		43
<i>ciMahya</i>	24	29	28	31	26	30	24	




Fig. 2 Expression profile of honeybee *Mahya*. *Mahya* and *EF-1alpha* mRNA levels are analyzed in the brains of newly emerged bees (day 0), 7-day-old in-hive bees, and 28-day-old foragers by Northern blot (a). The expression of *Mahya* and *beta-actin* mRNAs in the abdomens (a), brains (b), and thoraxes (t) of workers and the brains of workers (W), males (M), and queens (Q) was analyzed by RT-PCR. These bands are not detected without reverse transcription. The numbers at the right side are molecular weight markers (b)

the possible N-terminal signal sequence, Kazal-type serine/threonine protease inhibitor domain, EF-hand calcium-binding domain, two immunoglobulin domains, and the C-terminal novel domain. We also identified mouse (*mMahya-1* and *mMahya-2*), human (*hMahya-1* and *hMahya-2*), zebra fish (*drMahya-1* and *drMahya-2*), and ascidian (*ciMahya*) orthologues of *Mahya* by BLAST search. All of the vertebrate *Mahya* and *ciMahya* proteins share the same domain structures as the honeybee *Mahya* (Fig. 1a). The N-terminal halves with known domains and the novel C-terminal halves of *Mahya* proteins are equally conserved across the species (Tables 1 and 2), indicating that the novel C-terminal domain is also important for the function of *Mahya*. We therefore name this domain as the *Mahya* domain. The phylogenetic relationship among *Mahya* proteins from different species (Fig. 1b) shows that the vertebrate *Mahya-1* and *Mahya-2* proteins cluster into the separate clades. Honeybee and ascidian *Mahya* are quite diverged from each other and from the vertebrate *Mahya*. The search for *Mahya* in the honeybee and ascidian genomes demonstrates that both species contain a single *Mahya* gene in the genome.




Fig. 3 Expression of honeybee *Mahya* in the brain. The expression areas of *Mahya* in the brain were analyzed by in situ hybridization of the antisense and sense riboprobes to the cryosections of honeybee brain. Frontal sections of the entire brain area (a), mushroom bodies (b), and antennal lobes (c) displaying specific expression of *Mahya* are shown. The black arrows indicate the Kenyon cells in the calyces of mushroom bodies, the white arrow indicates the neurons at the inner border of the calyces, and the black arrowheads point to the lateral neurons surrounding the antennal lobes in (a). The black arrows in (b) show the small cell-body Kenyon cells expressing the high level of *Mahya* mRNA. The above signals are absent in the section hybridized with the sense probes (d)

Meanwhile, two different *Mahya* genes are present in mouse and human genomes. These results suggest that the ancestral *Mahya* gene was duplicated once during the evolution of the vertebrates.

Mahya is absent in the genomes of fruit fly, mosquito, silk moth, and nematode

When we searched the genome databases for *Mahya* orthologues by BLASTP and TBLASTN, we were able to find them in *Strongylocentrotus purpuratus* (sea urchin), *Takifugu rubripes* and *Tetraodon nigroviridis* (teleost fish), *Xenopus tropicalis* (frog), *Gallus gallus* (chicken), *Rattus norvegicus* (rat), and *Pan troglodytes* (chimpanzee) genomes in addition to ascidian, zebra fish, mouse, and human. In contrast, we could not find the *Mahya* orthologues with the significant *E* values in the genomes of *Drosophila melanogaster*, *Anopheles gambiae*, *Bombyx mori*, and *Caenorhabditis elegans*. The exons encoding the Kazal-type serine/threonine protease inhibitor domain, EF-hand calcium-binding domain, and immunoglobulin domain are, of course, present in various genes of the above animals. However, the genome DNA sequences capable of encoding the *Mahya* domain cannot be identified by TBLASTN search with any *Mahya* proteins as a query. Honeybee belongs to Hymenoptera, and both fruit fly and mosquito are diptera. All are holometabolous insect orders with a similarly structured CNS. Honeybee and fruit fly are thought to have separated 280 million years ago (Carpenter and Burnham 1985). These results therefore suggest that the last common ancestors of Bilateria had the ancestral gene of *Mahya*, but it was lost or highly diverged in most of Protostome branches during evolution.

Expression profile of *Mahya* in honeybee

In contrast to *EF-1alpha* mRNA, *Mahya* mRNA level is higher in the brains of 7-day-old in-hive bees and 28-day-old foragers than newly emerged bees (Fig. 2a). Thus, *Mahya* is one of the up-regulated genes in the brains of honeybee workers by age or task. *Mahya* mRNA is detected in the brain but not in the thorax and abdomen of workers (Fig. 2b), suggesting that *Mahya* is expressed at high level in the brain compared to thoracic and abdominal ganglia. No caste-specific expression of *Mahya* is observed (Fig. 2b). By in situ hybridization, we found that *Mahya* is predominantly expressed in the antennal lobes and mushroom bodies, which are the centers of learning and memory in insect brains (Fig. 3). In the antennal lobes, *Mahya* mRNA is detected in the lateral and medial cells surrounding the antennal lobes (Fig. 3c). They make synaptic contacts with the glomerular neuropils, where the sensory input from antennae is integrated. *Mahya* is also expressed at high level in the intrinsic Kenyon cells within the calyces of mushroom bodies and the neurons at the inner border of the calyces in the posterior protocerebrum (Fig. 3a). It is likely that these neurons are a different population of Kenyon cells. The level of *Mahya* mRNA is higher in the small cell-body Kenyon cells located in the center of each of the calyces of mushroom bodies than in the large cell-body Kenyon cells (Fig. 3b) located within the calyces surrounding the central smaller Kenyon cells.




Fig. 4 Expression of zebra fish *Mahya* orthologues in the brain. The expression areas of *drMahya-1* (a–e) and *drMahya-2* (f–j) in the brain were analyzed by in situ hybridization of the antisense and sense oligonucleotide probes to the cryosections of zebra fish brain. The sagittal sections of the entire brain areas hybridized with the antisense probes are shown (a, f). The high level of *drMahya-1* mRNA is detected in the dorsal telencephalon (the white arrowhead in b) and the ventral corpus cerebelli (the white arrowhead in d). These signals are absent in the sections hybridized with the sense probe (c, e). The high level of *drMahya-2* mRNA is detected in the habenula (the white arrowhead in g). The intermediate level of the expression is also observed in the olfactory bulb (the white arrow in g) and the optic tectum (the white arrowhead in i). These signals are absent in the sections hybridized with the sense probe (h, j)

Most of the small Kenyon cells project to the basal ring of the mushroom body, and the large Kenyon cells project to the lip and collar region of the mushroom body.

Expression areas of *drMahya-1* and *drMahya-2* in the zebra fish brain

A high level of *drMahya-1* mRNA is detected in the dorsal telencephalon and ventral corpus cerebelli of zebra fish brain (Fig. 4a,b,d). Meanwhile, the expression of *drMahya-2* mRNA is very high in the habenula and moderately high in the olfactory bulb and optic tectum (Fig. 4f,g,i).

Expression areas of *mMahya-1* and *mMahya-2* in the mouse brain

The level of *mMahya-1* mRNA is particularly high in the olfactory bulb, hippocampus, and cerebellum of mouse brain (Fig. 5a). The detailed analysis at high magnification demonstrates that *mMahya-1* mRNA is expressed in the granular layer of the cerebellum (Fig. 5c), the hippocampal CA3 region (Fig. 5f), and the glomerular layer and mitral cell layer of the olfactory bulb (Fig. 5i). Meanwhile, the expression level of *mMahya-2* is lower than that of *mMahya-1*, although it is expressed in the hippocampus (Fig. 5k) and weakly in the cerebellum and olfactory bulb (Fig. 5m,s). The observation at high magnification indicates that *mMahya-2* mRNA is expressed in the CA3 pyramidal

cells and dentate gyrus granular cells of hippocampus (Fig. 5p).

Analysis of the existence of *Mahya* in the genomes of bee species of Apoidea

All five exons encoding the *Mahya* domain are present in both *Apis mellifera* and *A. cerana* (Fig. 6a), suggesting that *A. cerana* contains *Mahya*. Three stingless bees, *Nannotrigona perilampoides*, *Trigona guianae*, and *Tetragonula fuscobalteata*, are positive for all of the five exons except exon 14 in *Nannotrigona perilampoides* (Fig. 6b). These results demonstrate that the highly social stingless bees contain *Mahya*. The five exons are also present in four bumblebee species, *Bombus diversus*, *B. griseocollis*, *B. bimaculatus*, and *B. transversalis* (Fig. 6c). The bumblebees with intermediate sociality have *Mahya*. The mostly solitary orchid bee, *Eulaema meriana*, is positive for all five exons, although the efficiency of PCR amplification is low, probably due to sequence mismatches (Fig. 6b). Nevertheless, this solitary bee contains *Mahya*. The same results were obtained with the communal carpenter bees, *Xylocopa virginica* and *X. appendiculata* (Fig. 6d), demonstrating that they have *Mahya*. We then analyzed the existence of




Fig. 5 Expression of mouse *Mahya* orthologues in the brain. The expression areas of *mMahya-1* (a–j) and *mMahya-2* (k–t) in the brain were analyzed by in situ hybridization of the antisense and sense oligonucleotide probes to the cryosections of mouse brain. The sagittal sections of the entire brain areas hybridized with the antisense probes are shown (a, k). The Nissl staining patterns (b, e, h, l, o, r) and autoradiographs are shown. The high level of *mMahya-1* mRNA is detected in the granular layer of the cerebellum

(c), hippocampal CA3 region (f), and the glomerular and mitral cell layers of the olfactory bulb (i). These signals are absent in the sections hybridized with the sense probe (d, g, j). The high level of *mMahya-2* mRNA is detected in the hippocampal CA3 region and dentate gyrus granular cell layer (p). The low level of the expression is also observed in the granular layer of the cerebellum (m) and the mitral cell layer of the olfactory bulb (s). These signals are absent in the sections hybridized with the sense probe (n, q, t)




Fig. 6 The existence of *Mahya* in the genomes of various bee species of Apoidea. The presence of exon sequences encoding the *Mahya* domain (exons 13–17) in the genomes of different bee species was analyzed by genomic PCR. The results of honeybees (a, *Apis mellifera* and *A. cerana*), orchid bee (b, *Eulaema meriana*), stingless bees (b, *Nannotrigona perilampoides*, *Trigona guianae*, and *Tetragonula fuscobalteata*), bumblebees (c, *Bombus diversus*, *B. griseocollis*, *B. bimaculatus*, and *B. transversalis*), and carpenter bees (d, *Xylocopa virginica* and *X. appendiculata*) are shown. All of the

above bees belong to Apidae. The results of four different Apoidea bee species other than Apidae (e, *Andrena tsukubana*, *Lasioglossum occidens*, *Megachile tsurugensis*, and *Colletes patellatus*) are also shown. Three bands indicated by the white asterisks in *X. appendiculata* and *Andrena tsukubana* are nonspecifically amplified products that are not related to the *Mahya* domain encoding exons. The numbers at the left sides of the panels are molecular weight markers. All of PCR products were sequenced to confirm their identity as exons encoding the *Mahya* domain

Mahya in the genomes of bees belonging to the broader Apoidea. *Andrena tsukubana* is positive for all of the five exons, *Megachile tsurugensis* is positive for exons 13, 15, and 17, *Lasioglossum occidens* is positive for only exon 17, and *Colletes patellatus* is positive only for exon 15 (Fig. 6e). Thus, *Andrena tsukubana* and *Megachile tsurugensis* are likely to contain *Mahya*. *Lasioglossum occidens* and *Colletes patellatus* also may have *Mahya* since none of the *Mahya* domain-encoding exons is present in the genomes of fruit fly, mosquito, silk moth, and nematode. All of PCR products were sequenced to confirm that they correspond to exons encoding the *Mahya* domain.

Discussion

Functional implication of *Mahya* genes

Based on the amino acid sequences of *Mahya* proteins from all species examined, they appear to be secretory

proteins with several function-known domains (Fig. 1a). In fact, a fraction of mMaha-1 protein can be detected in the culture medium when it is expressed in HEK293 cells (not shown). The N-terminal domain structure of *Mahya* with a Kazal-type serine/threonine protease inhibitor domain and EF-hand calcium-binding domain is also found in follistatin and follistatin-like proteins that bind the TGF-beta family to inhibit their activities (Balemans and Hul 2002). The TGF-beta family exerts its diverse functions by binding its specific receptors to activate the cell-signaling pathways mediated by the Smad family (Shi and Massague 2003). Several TGF-beta proteins are expressed in many brain areas including the hippocampus and are known to stimulate dendrite outgrowth and neuronal differentiation (Ishihara et al. 1994). In addition to the above domains, *Mahya* also contains two immunoglobulin domains that bind a wide variety of proteins. These results suggest that *Mahya* may act as a scaffold protein linking the TGF-beta family and other unknown proteins to modulate their functions in the brain. The C-

terminal novel Mahya domain has a weak homology to a beta chain of quinoxinase that forms a seven-bladed beta-propeller and is a part of the enzyme active site (Datta et al. 2001). The Mahya domain may fold into a similar structure.

According to the expression patterns of *Mahya* genes in honeybee, zebra fish, mouse, and human, *Mahya* appears to have major functions in the brain compared to other nervous systems and tissues (Fig. 2). The antennal lobes of the honeybee brain and the olfactory bulb of the vertebrate brain, where *Mahya* is expressed, share the same functions: the integration and processing of olfactory information by the projection of olfactory neurons. Honeybee and fruit fly have structurally similar antennal lobes, but the expression of *Mahya* in honeybee and not fruit fly neurons (Fig. 3c) indicates the mechanistic differences of the integration and processing of olfactory cues. Honeybee *Mahya* is also expressed in the Kenyon cells of the mushroom bodies (Fig. 3b). The dendrites of the Kenyon cells arborize calyces, where the olfactory and visual inputs from projection neurons of the antennal and optic lobes are integrated. The axons of Kenyon cells bifurcate into two lobes, where the efferent connections with other protocerebral neurons are made. Mushroom bodies are thus thought to be higher-order neuronal structures engaged in multisensory integration, learning, and memory in insects. The basic structures of the mushroom bodies are also similar between honeybee and fruit fly, suggesting that *Mahya* is related to the functional but not gross structural differences between honeybee and fruit fly Kenyon cells.

The optic tectum of teleost fish where *Mahya* is expressed (Fig. 4i) is the integration area of visual and other sensory inputs. Because the optic tectum also projects to many other brain regions, it is the brain area of connecting various sensory input and motor output (Yoshimoto and Ito 1993). The habenula, another *Mahya*-expressing brain area (Fig. 4g), was shown to integrate photic inputs from the pineal as well as other brain regions and also project to many other brain areas (Yanez and Anadon 1996). However, its precise functions have not been identified yet. The dorsal region of the teleost fish telencephalon also expresses *Mahya* (Fig. 4b) and functions in the processing of multiple sensory inputs (olfactory, auditory, mechanosensory, and visual; Prechtl et al. 1998). The recent study with goldfish (Portavella et al. 2004) also demonstrates that the medial and lateral telencephalic regions are involved in the emotional memory system and the spatial and relational memory system, respectively. Thus, the medial and lateral telencephalic regions appear to correspond to the mammalian amygdala and hippocampus, respectively. The mammalian hippocampus is involved in spatial learning, relational memory, and processing of temporal attributes of events and situations. Both *mMahya-1* and *mMahya-2* are expressed in the CA3 region and the dentate gyrus of the hippocampus (Fig. 5). The vertebrate cerebellum, where *drMahya-1* and *mMahya-1* are highly expressed, functions in motor coordination and motor learning and memory. Moreover, the

cerebellum may be involved in emotional and fear memory (Yoshida et al. 2004; Sacchetti et al. 2002). The expression of human *Mahya* (*hMahya-1* and *hMahya-2*) mRNAs is also highest in the brain compared to the other tissues. In the human brain, both *hMahya-1* and *hMahya-2* mRNAs are detected in most of areas; however, *hMahya-1* mRNA is relatively high in the cerebellum and amygdala, and *hMahya-2* mRNA is high in the amygdala and thalamus (Kikuno et al. 2004; <http://www.kazusa.or.jp/huge/gfpage/KIAA1263/> for *hMahya-1* and <http://www.kazusa.or.jp/huge/gfpage/KIAA1061/> for *hMahya-2*). The thalamus integrates all sensory information except olfactory and transmits these information back and forth between the cortex. Thus, *Mahya* genes are expressed in the functionally equivalent brain areas of the honeybee, zebra fish, mouse, and human, suggesting that the basic function of *Mahya* should be conserved in the brains of honeybee and vertebrates.

Mahya appears to be present in diverse bees classified as Apoidea (Fig. 6). The possibility that the *Mahya* genes we detected in the various bees by genomic PCR are pseudogenes could not be completely ruled out at this stage (it is, of course, necessary to isolate and sequence the full-length cDNAs). Nevertheless, we think this is unlikely because both *Drosophila melanogaster* and *Anopheles gambiae* genomes were shown to have small numbers of pseudogenes (176 and 166, respectively; Zdobnov et al. 2002) compared to mammals (more than 10,000). Also, the *Mahya* domain encoding exon sequence is absent in these model insects and silk moth genome. The bees apparently containing *Mahya* include both solitary and highly social bees. However, these bee species construct nests and bring food back to their nest for storage and for raising progeny. These results thus suggest that *Mahya* is not related to the high degree of sociality (colony maintenance by multiple individuals, caste differentiation, and generation overlap) but may be involved in the cognitive ability of bees associated with accurate spatial and olfactory recognition and memory (Giurfa et al. 2001).

What is the function of *Mahya* in sea urchin and sea squirt with relatively simple nervous systems, which do not exhibit complex behavior? If *Mahya* genes are expressed in the nervous systems of above animals, they would be involved in the primary processing of chemosensory information and not high order of learning and memory processes. Thus, the basic functions of *Mahya* conserved between Deuterostomes and Hymenoptera should be related with the mechanistic aspects of sensory information processing, which are different from those in fruit fly, mosquito, silk moth, and nematode. However, the possibility that *Mahya* genes are expressed in tissues other than nervous systems of sea urchin and sea squirt cannot be ruled out at this point.

The genome of Hymenoptera

It was reported that approximately 10% of the annotated genes in *Drosophila melanogaster* and *Anopheles gambiae*

show better sequence matches with noninsect genes (Zdobnov et al. 2002). Although the percentage of this category of genes will likely be reduced after the completion of the *Apis mellifera*, *Bombyx mori*, and *Tribolium castaneum* genome projects, each insect must contain genes with specific functions shared between noninsect animal species. The existence of these genes reflects the selection and subsequent adaptation associated with different ecologies and life histories. In the case of *Apis mellifera*, we could predict the genes related with the high degree of sociality and cognitive ability. Furthermore, we can also expect honeybee-specific genes involved in polyphenism (caste differentiation) and in sex determination via haplodiploidy.

Gene loss and divergence in the evolution of metazoan genomes

Distribution of *Mahya* across Bilaterian species and phylogenetic relationship among *Mahya* proteins from different species suggest that the last common ancestors of Bilateria (Urbilateria) had the ancestral *Mahya* genes. It also indicates that Urbilateria was genetically quite complex with diverse sets of genes. We propose that the ancestral gene of *Mahya* was lost or became highly diverged in many metazoan species other than Hymenoptera and Deuterostomes. Even if highly diverged *Mahya* is present in many metazoan genomes, the ancestral function of *Mahya* would be more conserved between Hymenoptera and Deuterostomes. In terms of searching for the true ancestor of *Mahya*, it will be interesting to see if *Mahya* is present in the genomes of Cnidaria. This type of gene loss is now recognized as one of the scenarios leading to metazoan diversity (Kortschak et al. 2003). Even between two dipteran insects, fruit fly and mosquito, that diverged 250 million years ago, only mosquito contains genes homologous to human leukotriene B4 12-hydroxy dehydrogenase and calcineurin-binding intracellular regulatory protein (Zdobnov et al. 2002). Since fruit fly and honeybee diverged about 280 million years ago (Carpenter and Burnham 1985), we could also expect honeybee-specific gene loss and retention. Cross-species genome sequence comparison and gene functional analysis will undoubtedly give insights into how gene emergence, divergence, and loss have led the spectacular diversity of metazoans.

Acknowledgements We thank M. Mizunami for helpful discussion. This study was supported by Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science to T.K.

References

Ashburner M (1989) *Drosophila a laboratory manual*. Cold Spring Harbor Laboratory Press, New York

Balemans W, Hul WV (2002) Extracellular regulation of BMP signaling in vertebrates: a cocktail of modulators. *Dev Biol* 250:231–250

Carpenter FM, Burnham L (1985) The geological record of insects. *Annu Rev Earth Planet Sci* 13:297–314

Datta S, Mori Y, Takagi K, Kawaguchi K, Chen ZW, Okajima T, Kuroda S, Ikeda T, Kano K, Tanizawa K, Mathews FS (2001) Structure of a quinoxaline amine dehydrogenase with an uncommon redox cofactor and highly unusual crosslinking. *Proc Natl Acad Sci U S A* 98:14268–14273

Fahrbach SE, Robinson GE (1995) Behavioral development in the honey bee: toward the study of learning under natural conditions. *Learn Mem* 2:199–224

Funada M, Yasuo S, Yoshimura T, Ebihara S, Sasagawa H, Kitagawa Y, Kadowaki T (2004) Characterization of the two distinct subtypes of metabotropic glutamate receptor from honeybee, *Apis mellifera*. *Neurosci Lett* 359:190–194

Giurfa M, Zhang S, Jenett A, Menzel R, Srinivasan MV (2001) The concepts of ‘sameness’ and ‘difference’ in an insect. *Nature* 410:930–933

Ishihara A, Saito H, Abe K (1994) Transforming growth factor-beta 1 and -beta 2 promote neurite sprouting and elongation of cultured rat hippocampal neurons. *Brain Res* 639:21–25

Kikuno R, Nagase T, Nakayama M, Koga H, Okazaki N, Nakajima D, Ohara O (2004) HUGE: a database for human KIAA proteins, a 2004 update integrating HUGEppi and ROUGE. *Nucleic Acids Res* 32(database issue):D502–D504

Kortschak RD, Samuel G, Saint R, Miller DJ (2003) EST analysis of the Cnidarian *Acropora millepora* reveals extensive gene loss and rapid sequence divergence in the model invertebrates. *Curr Biol* 13:2190–2195

Menzel R, Giurfa M (2001) Cognitive architecture of a mini-brain: the honeybee. *Trends Cogn Sci* 5:62–71

Menzel R, Muller U (1996) Learning and memory in honeybees: from behavior to neural substrates. *Annu Rev Neurosci* 19:379–404

Mineta K, Nakazawa M, Cebria F, Ikeo K, Agata K, Gojobori T (2003) Origin and evolutionary process of the CNS elucidated by comparative genomics analysis of planarian ESTs. *Proc Natl Acad Sci U S A* 100:7666–7671

Mushegian AR, Garey JR, Martin J, Liu LX (1998) Large-scale taxonomic profiling of eukaryotic model organisms: a comparison of orthologous proteins encoded by the human, fly, nematode, and yeast genomes. *Genome Res* 8:590–598

Portavella M, Torres B, Salas C (2004) Avoidance response in gold fish: emotional and temporal involvement of medial and lateral telencephalic pallium. *J Neurosci* 24:2335–2342

Prechtl JC, von der Emde G, Wolfart J, Karamursel S, Akoev GN, Andrianov YN, Bullock TH (1998) Sensory processing in the pallium of a mormyrid fish. *J Neurosci* 18:7381–7393

Sacchetti B, Baldi E, Lorenzini CA, Bucherelli C (2002) Cerebellar role in fear-conditioning consolidation. *Proc Natl Acad Sci U S A* 99:8406–8411

Shi Y, Massague J (2003) Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* 113:685–700

Tsuchimoto M, Aoki M, Takada M, Kanou Y, Sasagawa H, Kitagawa Y, Kadowaki T (2004) The changes of gene expression in honeybee (*Apis mellifera*) brains associated with ages. *Zool Sci* 21:23–28

Whitfield CW, Brand MR, Bonaldo MF, Kumar CG, Liu L, Pardinas JR, Robertson HM, Soares MB, Robinson GE (2002) Annotated expressed sequence tags and cDNA microarrays for studies of brain and behavior in the honey bee. *Genome Res* 12:555–566

Yanez J, Anadon R (1996) Afferent and efferent connections of the habenula in the rainbow trout (*Oncorhynchus mykiss*): an indocarbocyanine dye (DiI) study. *J Comp Neurol* 372:529–543

- Yoshida M, Okamura I, Uematsu K (2004) Involvement of the cerebellum in classical fear conditioning in gold fish. *Behav Brain Res* 153:143–148
- Yoshimoto M, Ito H (1993) Cytoarchitecture, fiber connections, and ultrastructure of the nucleus pretectalis superficialis pars magnocellularis (PSm) in carp. *J Comp Neurol* 336:433–446
- Yoshimura T, Suzuki Y, Makino E, Suzuki T, Kuroiwa A, Matsuda Y, Namikawa T, Ebihara S (2000) Molecular analysis of avian circadian clock genes. *Brain Res Mol Brain Res* 78:207–215
- Zdobnov EM, von Mering C, Letunic I, Torrents D, Suyama M, Copley RR, Christophides GK, Thomasova D, Holt RA, Subramanian GM, Mueller H-M, Dimopoulos G, Law JH, Wells MA, Birney E, Charlab R, Halpern AL, Kokoza E, Kraft CL, Lai Z, Lewis S, Louis C, Barillas-Mury C, Nusskern D, Rubin GM, Salzberg SL, Sutton GG, Topalis P, Wides R, Wincker P, Yandell M, Collins FH, Ribeiro J, Gelbart WM, Kafatos FC, Bork P (2002) Comparative genome and proteome analysis of *Anopheles gambiae* and *Drosophila melanogaster*. *Science* 298:149–159