

Adaptive evolution of color vision of the Comoran coelacanth (*Latimeria chalumnae*)

SHOZO YOKOYAMA*, HUAN ZHANG†, F. BERNHARD RADLWIMMER, AND NATHAN S. BLOW

Department of Biology, Syracuse University, Syracuse, NY 13244

Communicated by M. T. Clegg, University of California, Riverside, CA, March 26, 1999 (received for review December 5, 1998)

ABSTRACT The coelacanth, a “living fossil,” lives near the coast of the Comoros archipelago in the Indian Ocean. Living at a depth of about 200 m, the Comoran coelacanth receives only a narrow range of light, at about 480 nm. To detect the entire range of “color” at this depth, the coelacanth appears to use only two closely related paralogous RH1 and RH2 visual pigments with the optimum light sensitivities (λ_{\max}) at 478 nm and 485 nm, respectively. The λ_{\max} values are shifted about 20 nm toward blue compared with those of the corresponding orthologous pigments. Mutagenesis experiments show that each of these coadapted changes is fully explained by two amino acid replacements.

The coelacanth, *Latimeria chalumnae*, is the sole surviving species of the old lineage of crossopterygian fishes that were abundant during the Devonian period. The morphology of the coelacanth has changed very little over 400 million years (1–3). This “living fossil” lives at a depth of about 200 m near the Comoros archipelago in the western Indian Ocean (4, 5). Living at a depth of 200 m, the Comoran coelacanth receives only a narrow range of color, at about 480 nm (6). To see in this photic environment, the retina of the coelacanth consists mostly of rods, with a much smaller number of cones (7, 8). The molecular genetic basis of coelacanth vision is unknown, although one visual pigment from the Comoran coelacanth has been identified (9). Here we report the structure and function of the opsin genes of the coelacanth and elucidate how the coelacanth has modified its visual pigments to adapt to its natural habitat.

Visual pigments consist of a chromophore, 11-*cis* retinal, and a transmembrane protein, an opsin (10). Vertebrate retinal opsins (or pigments) are classified into five evolutionarily distinct groups: (i) rhodopsin (RH1), (ii) RH1-like (RH2), (iii) short wavelength-sensitive (SWS1), (iv) SWS1-like (SWS2), and (v) long wavelength-sensitive (LWS) or middle wavelength-sensitive (MWS) (LWS/MWS) opsin clusters (11–13). RH1 pigments are usually expressed in rods and the four other classes of pigments in cones. All five classes of pigments appear to have coexisted in the common ancestor of vertebrates (11–13). A central unanswered question in phototransduction is how these visual pigments achieve the wavelength of maximal absorption (λ_{\max}) ranging from 360 nm (ultraviolet) to 700 nm (infrared). Recent molecular genetic analyses show that the difference in the λ_{\max} values of LWS/MWS pigments in mammals is primarily controlled by amino acids at five critical sites (14). Receiving a restricted range of light, the Comoran coelacanth provides an excellent opportunity to study not only the molecular basis of blue-shifted color vision but also the processes of adaptive evolution to its unique photic environment.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 18 U.S.C. §1734 solely to indicate this fact.

PNAS is available online at www.pnas.org.

MATERIALS AND METHODS

Genomic DNA Library Cloning and DNA Sequencing. Genomic DNA was isolated from the Comoran coelacanth, which was obtained from the American Museum of Natural History, NY (tissue loan no. AMNH 59196). A genomic library of this coelacanth was constructed with *Bam*HI-digested λ EMBL3 vector and *Sau*3AI partially digested coelacanth genomic DNA (10–20 kb). By using bovine RH1 and human SWS1 cDNAs as probes, we screened about 1.4×10^6 and 10^7 recombinant plaques, respectively. From the first screening, we isolated two clones, λ Lc16 and λ Lc58, which represent two distinct functional opsin genes. From the second screening, we isolated two nonoverlapping clones, λ Lc17 and λ Lc01, which represent a pseudogene. These genomic clones were subcloned into pBluescript SK(–), and the nucleotide sequences were determined for both strands by the dideoxy-chain-termination method (15).

Creation of Minigenes and Site-Directed Mutagenesis. The five exon regions of each of the two functional opsin genes were PCR-amplified by using five sets of oligonucleotides. For λ Lc16, we used 5'-CCAAGGGAATTCCACCATGAATGGAACAGAGGGTCC-3' (forward) and 5'-TTACCTCCTAGGGTAGCGAAAAATCCCTCAAT-3' (reverse) for exon 1, 5'-CCTTCCCCTAGGAGGTCAAGTTGCTCTGTGG-3' (forward) and 5'-GAGCTATCTAGACCATCCGAAAAGAGGAGGAAC-3' (reverse) for exon 2, 5'-CTGAATTCTAGATATATCCAGAGGGTATGCA-3' (forward) and 5'-ACAAAGCTGCAGCATCCTTGACAGTGCAGAC-3' (reverse) for exon 3, 5'-TTAGGCTGCAGCAGCAGCAGGAGTCTGCC-3' (forward) and 5'-TTTG-CATCCGGA~~ACTGTTT~~GTTTCAGGAGAATGTAG-3' (reverse) for exon 4, and 5'-TACAGTCCGGA~~ACTGCATGATCACCACCCTC~~-3' (forward) and 5'-CCGAGGTCGACGCGGGTGAGACACTGCTGGA-3' (reverse) for exon 5. The underlined sequences indicate restriction enzyme site for cloning purposes. The extra 5–6 nt at the 5' end are to facilitate restriction enzyme digestion. For λ Lc58, we used 5'-CAAACAGAATTCACCATGAACGGCAGCGAA-3' (forward) and 5'-CTACCTCCTAGGGTGGCAAAGAAACCTCCAT-3' (reverse) for exon 1, 5'-TTCCTCCTAGGAGGTCAGGTAGCTCTCTGG-3' (forward) and 5'-ACCAA TCTAGACCAGCCAACGAGTGGCGGTGC-3' (reverse) for exon 2, 5'-CCTTTTCTAGATATATC-CCCAGGGACTGCAG-3' (forward) and 5'-AAACA~~ACTCGAGCCTCTT~~GACTTTGCAGAT-3' (reverse) for exon 3, 5'-CCAGGCTGCAGCACAAACAGCAGGAATCTGCA-3'

Abbreviations: RH1_{Lc}, rhodopsin (or pigment) of coelacanth; RH2_{Lc}, RH1-like opsin (or pigment) of coelacanth; SWS, short wavelength-sensitive; MWS, middle wavelength-sensitive; LWS, long wavelength-sensitive.

Data deposition: The sequences reported in this paper have been deposited in the GenBank database (accession nos. AF131253–AF131257 for *rh1*_{Lc} and AF131258–AF131262 for *rh2*_{Lc}).

*To whom reprint requests should be addressed. e-mail: syokoyam@mailbox.syr.edu.

†Present address: Irigo Institute, Atsumi-cho, Atsumi-gun, Aichi-ken, Japan.




FIG. 1. λ clones and sequencing strategies of $rh1_{Lc}$ and $rh2_{Lc}$. The five exons are indicated by black boxes in the restriction maps. DNA segments subcloned and sequenced are represented as thick bars in λ clones. A, *Sall*; B, *Bam*HI; H, *Hind*III; K, *Kpn*I; S, *Sst*I.

(forward) and 5'-AAGAGCATCCGGAAGCTTGT-TCAACAACAC-3' (reverse) for exon 4, and 5'-TCTAGT-TCCGGAAGCTTGTATGATCACCACGTTG-3' (forward) and 5'-GAAGTGGTCGACGCGGGCGAAACCTGACTGG-A-3' (reverse) for exon 5. PCR was performed by using 30 cycles at 92°C for 45 sec, 55°C for 60 sec, and 72°C for 90 sec. At each cycle, the duration of the extension reaction was progressively extended by 3 sec. The PCR products were digested by appropriate restriction enzymes and ligated into pBluescript SK (-). Nucleotide sequences of the entire regions of the minigenes were determined. We selected clones that

encode identical amino acid sequences to those of the corresponding sequences deduced from the genomic sequences.

Mutants were generated by using QuikChange site-directed mutagenesis kit from Stratagene. All DNA fragments that were subjected to mutagenesis were sequenced to rule out spurious mutations.

Absorption Spectra. The two coelacanth minigenes were ligated into an expression vector pMT, expressed in cultured COS1 cells, and regenerated with 11-*cis* retinal, and the resulting visual pigments were purified as described (16, 17). UV-visible spectra were recorded at 20°C by using a Hitachi U-3000 dual beam spectrophotometer. Visual pigments were bleached for 3 min by using a 60 W standard light bulb equipped with a Kodak Wratten no. 3 filter at a distance of 20 cm. Data were analyzed with SIGMAPLOT software (Jandel, San Rafael, CA).

Sequence Data Analyses. The amino acid sequences of the coelacanth visual pigments were compared with those of other vertebrate pigments. The root of the phylogenetic tree of these pigments was determined by using goldfish LWS (GenBank accession no. L11867), African clawed frog LWS (U90895), chicken LWS (M62903), and human MWS (K03490-K03497) pigments as the outgroup. The number (K) of amino acid replacements per site for a pair of sequences was estimated by $K = -\ln(1 - p)$, where p is the proportion of different amino acids per site. Topology and branch lengths of the phylogenetic tree were evaluated by using the neighbor-joining method (18). The reliability of the neighbor-joining tree topology was evaluated by the bootstrap analysis with 1,000 replications (19).

The amino acid sequences of ancestral pigments were inferred by using likelihood-based Bayesian method (20) with a modified version of the Jones, Taylor, and Thornton (JTT; ref.

		I	
Coelacanth RH1	MNGTEGPNFYVPM SNKTVVRNPF GYPQYYLADPWKYSALAA YMFFLILVGFPI NFLT L FVTIQHKKLRTP LNYILL		77
Coelacanth RH2	MNGTEGMNFYVPLSNRTGLVRS PF EY TQYYLAE PWKFSVLCAYMFL LIILGFPINFLTLLVTFKHKKLRQPLNYILV		77
Coelacanth ψ B	M. .LRGEDFYL. FSKISSV. .GLWDGLQYHIAP I WAFY LQAVFMGF MFFVGTLLNAIVLIVTLKY NKL*QPLNYILL		72
		tga	
	II	III	IV
RH1	DLAVADLCMVFGGFFVTMYSSMNGYFVLGPTGCNIEGFFATLGGQVALWALVVLAIERYVVVCKPMSNFRFGENHAIMGVIFTWIMAL		165
RH2	NLAVASLFMVVFGFTVTFYSSLNGYFVLGPMGCAMEGFFATLGGQVALWLSLVLA IERYIVVCKPMGNFRFASHAIMGIAFTWIMAL		165
ψ B	NISLAGFIVCLF. IFTVFISSRGYFIFGRAVCAIESFLGSM TGLITG. . . TLAFERDIVICKPFSNFCFGSKQSL LAVGATWVIGL		160
	c	a	
		V	
RH1	SCAVPPLFGWSRYIPEGMQSSCGVDYITLKP E INNESFVIYMFVVHFTIPLIVIFFCYGRVCTVKD. .AAAQQQESATTQKAKEV T		251
RH2	ACAAPPLVGWSRYIPEGLQCS CGPDYITLNPDFHNESYVMYLFVHFLLPITTIFFTYGR LICKVKE. .AAAQQQESASTQKAKEV T		251
ψ B	GAALPPVFGWSQYIPEGLQCS GP EWYTVNKNWNNESYVIFLFSFCFGLPLSIIIFS YTKLLM TLHT. *VAKQQEQSASAQKAEREV T		248
		a taa	
	VI	VII	
RH1	RMVIVMVISFLVCWVPYASVAGYIFFNQSEFGPVFMTAPSF FAKSASFYNPVIYFLLNKQFRNCMITTLCCGKNPFGDE DATSSAGS		339
RH2	RMVILMVIGFLTAWVPYASAAF WIFCNRGEEFTATLMTVPAFFSKSSCLFNPIIYVLLNKQFRNCMITRLCCGENPLGDDDTSSAVSQ		339
ψ B	KMVIVMVLGFLMCWLPYASFALWVITHRGEPFDLRMASIPSVFSKASTVYNPIIYIFMKNQ.		310
RH1	SKTEASSVSSSSVSPA		355
RH2	RKTDVSSVSSQLSPA		355
ψ B		

FIG. 2. The deduced amino acid sequences of coelacanth RH1 (RH1_{Lc}), coelacanth RH2 (RH2_{Lc}), and coelacanth ψ B (ψ B_{Lc}). Gaps, denoted by dots, were inserted in the sequences for the optimal alignment. Deletion, insertion, and nonsense nucleotide mutations in ψ sws1_{Lc} are noted by lower case letters. * indicates a stop codon. Exon 5 of ψ sws1_{Lc} is missing. Putative transmembrane domains I-VII are indicated by horizontal lines.




FIG. 3. Phylogenetic tree of vertebrate visual pigments whose λ_{max} values (given after P) have been evaluated. Opsin (or visual pigment) names and their references (accession numbers) are: lamprey, *Lamptera japonica* (M63632); mosquito fish, *Gambusia affinis* (Y11146); goldfish, *Carassius auratus* (L11863); cavefish, *Astyanax fasciatus* (U12328); African clawed frog, *Xenopus laevis* (L07770); bullfrog, *Rana catesbeiana* (S79840); leopard frog, *Rana pipiens* (S49004); salamander, *Ambystoma tigrinum* (U36574); American chameleon, *Anolis calominensis* (L31503); chicken, *Gallus gallus* (D00702); bovine, *Bos taurus* (M21606); and dolphin, *Tursiops truncatus* (AF055456) for RH1 cluster; goldfish-G1 (L11865) and -G2 (L11866); chicken (M92038); American chameleon (31); and gecko, *Gekko gekko* (M92035) for RH2 cluster; goldfish (D85863); African clawed frog (U23463); American chameleon (32); chicken (M92039); human, *Homo sapiens* (M13295); mouse, *Mus musculus* (U49720); and rat, *Rattus norvegicus* (U63972) for SWS1 cluster; and goldfish (L11864); American chameleon (32); and chicken (M92037) for SWS2 cluster.

21), Dayhoff (22), and the equal-input models (20) of amino acid replacements.

RESULTS AND DISCUSSION

Coelacanth Opsin Genes. To determine what types of opsin genes the Comoran coelacanths possess, we hybridized Southern blots containing the coelacanth genomic DNA, digested with various restriction enzymes, to bovine RH1 (23), human SWS1 (24), and human LWS (24) opsin cDNAs separately. The Southern analysis revealed strongly hybridizing bands to the first two probes, but no band to the third probe (data not shown). Thus, the coelacanth has RH1 and SWS1-like genes but no LWS/MWS genes.

After genomic DNA library screening using bovine RH1 opsin cDNA as a probe, we isolated 12 positive phage clones. Among these, λ Lc16 and λ Lc58 were found to contain the

entire coding regions of functional opsin genes, designated as $rh1_{Lc}$ and $rh2_{Lc}$, respectively (Fig. 1). Similarly, by using human SWS1 opsin cDNA as a probe, we isolated 13 positive clones. Among these, 2 nonoverlapping clones, λ Lc17 containing exons 1 and 2 and λ Lc01 containing exons 3 and 4, were found to represent a pseudogene, designated as ψ sws1_{Lc}. As previously noted, the coelacanth does not have any LWS/MWS opsin genes. From an extensive screening of genomic DNA library with human SWS1 opsin cDNA as a probe, we could isolate only one SWS1 pseudogene. This strongly suggests that the coelacanth has also lost a functional SWS2 opsin gene.

The RH1_{Lc} and RH2_{Lc} opsins, deduced from the DNA sequences of $rh1_{Lc}$ and $rh2_{Lc}$, consist of 355 aa (Fig. 2). The pseudogene ψ sws1_{Lc} contains premature stop codons because of nonsense and frameshift mutations (Fig. 2). When the coding regions of $rh1_{Lc}$ and $rh2_{Lc}$ are compared, the proportion of identical nucleotides is 70%. When $rh1_{Lc}$ and $rh2_{Lc}$ are compared with bovine $rh1$, the sequence similarities are given by 76% and 71%, respectively. Thus, $rh1_{Lc}$ is more closely related to bovine $rh1$ gene than to $rh2_{Lc}$, suggesting that $rh1_{Lc}$ and $rh2_{Lc}$ were duplicated before the vertebrate radiation.

Evolution of Coelacanth Visual Pigments. Fig. 3 shows the rooted phylogenetic tree of RH1_{Lc}, RH2_{Lc}, ψ B_{Lc}, deduced from ψ sws1_{Lc}, and RH1, RH2, SWS1, and SWS2 pigments in other vertebrates whose λ_{max} values have been directly determined. The bootstrap values of forming RH1, SWS1, and SWS2 clusters are 99%, 100%, and 100%, respectively, and are highly reliable, but that for RH2 cluster is only 81%. Note that RH1 cluster includes a wide range of vertebrates, from lamprey to mammals, and furthermore, the bootstrap value of forming the cluster of RH1 and RH2 groups is 100%. Thus, RH1 and RH2 clusters consist of two sets of paralogous pigments. These observations show that RH1_{Lc}, RH2_{Lc}, and ψ B_{Lc} belong to the RH1, RH2, and SWS1 clusters, respectively.

Within the SWS1 cluster, ψ B_{Lc} is most closely related to mammalian opsins, but this "misclassification" is probably caused by the much faster evolutionary rate of ψ sws1_{Lc}, a characteristic of pseudogenes. Fig. 3 suggests that, in the RH1 cluster, American chameleon (P491) pigment is not closely related to chicken (P503) pigment. This unexpected tree topology arises because of an accelerated rate of amino acid replacements in the former pigment in the rodless American chameleon (13). In the RH2 cluster, the American chameleon (P495) pigment is more closely related to the chicken pigment than to the gecko pigment. This may be explained not only by the accelerated evolutionary rate of the gecko pigment but also by the decelerated rate of the chicken pigment. This can be seen by making pairwise comparisons of chicken (P508), gecko (P467), and American chameleon (P495) pigments with an appropriate outgroup. For example, although they are not statistically significant, chicken (P508) pigment has shorter branch lengths than gecko (P467) and American chameleon (P495) pigments (Table 1). The branch length, 0.149, for gecko (P467) pigment is significantly longer than 0.003 for American chameleon (P495) pigment (Table 1).

It is widely accepted that tetrapods originated from lobe-finned fish, which include the coelacanth (25). According to

Table 1. The numbers of amino acid replacements per site (K × 100) in the RH2 cluster

Branch (pigment)			
Chicken (P508)	Gecko (P467)	American chameleon (P495)	Outgroup (pigment)
3.1 ± 0.95	—	4.9 ± 1.21	Goldfish (P511)
7.8 ± 1.52	12.2 ± 1.93	—	Goldfish (P511)
—	14.9 ± 2.15*	3.0 ± 0.93*	Chicken (P508)

*, Significantly different at 1% level.




FIG. 4. Amino acid replacements at informative residues of RH1_{Lc} and RH2_{Lc} pigments. Those surrounded by a single rectangle denote a 1-aa replacement event. Ancestral amino acids were inferred by considering tree topology: (((lamprey [P500], ((goldfish [P492], cavefish [P505])), (RH1_{Lc}, (African clawed frog [P502], (chicken [P503], bovine [P500])))), ((goldfish [P511], goldfish [P506])), (RH2_{Lc}, ((gecko [P467], American chameleon [P495]), chicken [P508])))), (goldfish [P455], (chicken [P455], American chameleon [P437]))).

this scenario, RH1_{Lc} and RH2_{Lc} pigments should be more closely related to tetrapod pigments than to ray-finned fish pigments within each orthologous group. Although the opsin sequences alone are not sufficient to resolve the divergence of the coelacanths, tetrapods, and bony fishes (Fig. 3), there is other considerable evidence to support a common ancestor for tetrapods and coelacanth (25). But, if the coelacanth is more closely related to tetrapods than to ray-finned fishes, then amino acid replacements that occurred in the early stages of the coelacanth evolution are expected to be shared more often with tetrapods than with ray-finned fishes. Such shared amino acid replacements can be traced by inferring the amino acid sequences of the ancestral RH1 and RH2 pigments. Applying the JTT model of amino acid replacements, we inferred the amino acid sequences of all ancestral RH1 and RH2 pigments. Then, by comparing the amino acids of RH1_{Lc} and RH2_{Lc} pigments with those at the corresponding sites of the orthologous ancestral pigments, 20 shared amino acid replacements can be identified (Fig. 4). Among these, 13 favor the closer

relationship of the coelacanth and tetrapods, whereas 7 disfavor the relationship. Even when the coelacanth pigments are assumed to be more closely related to ray-finned fishes than to tetrapods, the ratio between the favorable and unfavorable amino acid replacements becomes 12:7. Similar ratios were obtained with the Dayhoff and equal-input models of amino acid replacements (data not shown). Although none of the ratios is significantly different from 1:1, they do suggest a closer relationship between the coelacanth and tetrapods. Of interest, the *rh1* genes of the coelacanth and tetrapods contain four introns at identical positions, whereas those of ray-finned fishes contain no introns, having lost them during evolution (26). These observations are compatible with the notion that the coelacanth is more closely related to tetrapods than to ray-finned fishes.

Absorption Spectra of RH1_{Lc} and RH2_{Lc} Pigments. The spectral sensitivity of visual pigments usually is evaluated directly by expressing opsin cDNAs in cultured cells (16, 17). However, because the retina of the coelacanth is not available




FIG. 5. Absorption spectra of the wild-type (wt) RH1_{Lc} and RH2_{Lc} and mutant pigments evaluated by the dark-light difference spectra.

to us, the 5'- and 3'-flanking regions and the four intron segments of *rh1_{Lc}* and *rh2_{Lc}* were deleted by using recombinant DNA techniques (see *Materials And Methods*). The resulting minigenes containing only the five exon sequences were ligated into an expression vector pMT, expressed in COS1 cells, and reconstituted with 11-*cis* retinal. The RH1_{Lc} and RH2_{Lc} pigments constructed in this way were shown to have λ_{\max} values of 485 ± 3 nm and 478 ± 1 nm, respectively (Fig. 5). The latter λ_{\max} of the cone-specific pigment corresponds to the previously described pigment with a λ_{\max} of 473 nm (9), suggesting that the coelacanth pigment described by Dartnall (9) might have been sampled from a rare cone photoreceptor cell.

With the exception of gecko (P467) pigment, most RH1 and RH2 pigments have λ_{\max} values of about 500 nm (Fig. 3). Compared with them, the λ_{\max} values of RH1_{Lc} and RH2_{Lc} pigments are about 20 nm blue-shifted. The fascinating part of the two λ_{\max} values (478 nm and 485 nm) is that they correspond to the narrow range of light around 480 nm in the Comoran coelacanth's natural photic environment (6). Thus, the RH1_{Lc} and RH2_{Lc} pigments have coevolved to detect two edges of the available light spectra so that the coelacanth can distinguish the entire range of "colors" available to them. Because the coelacanth retina has both rods and cones (7, 8), the finding of a rod-specific RH1_{Lc} pigment and at least one cone-specific pigment is expected. However, it was unexpected to find that the coelacanth uses RH2 pigment instead of SWS1 or SWS2 pigments for its blue vision. It has been demonstrated that color vision is possible with only rods and a single class of cones in cats. By using rods and blue-sensitive cones simultaneously, cats can distinguish wavelengths in the range 440–500 nm at twilight (27). Evidently, the coelacanth uses rod-specific RH1_{Lc} pigments and cone-specific RH2_{Lc} pigments to visualize the entire, albeit limited, range of color available to them.

Adaptive Mechanisms of RH1_{Lc} and RH2_{Lc} Pigments. How did RH1_{Lc} and RH2_{Lc} pigments achieve the blue shifts? It is known that amino acid changes E122Q and A292S in the bovine (P500) pigment shift the λ_{\max} value 20 nm (28) and 10 nm (29) toward blue, respectively. Within the RH1 cluster, these two amino acid replacements E122Q and A292S occurred in the coelacanth after its divergence from other organisms (Fig. 3, branch a). Our mutagenesis analysis shows that single mutants Q122E and S292A and a double mutant Q122E/S292A of RH1_{Lc} pigment have λ_{\max} values of 495 ± 2 nm, 493 ± 1 nm, and 511 ± 2 nm, respectively (Fig. 5). Thus, E122Q and A292S together can cause a 25 nm blue-shift in the λ_{\max} and fully explain the observed λ_{\max} of RH1_{Lc} pigment.

The blue-shifted λ_{\max} of RH2_{Lc} pigment also is explained by two amino acid replacements, E122Q and M207L. Q122 is shared by the RH2 pigments of coelacanth, gecko, chicken, and American chameleon, and the E122Q replacement occurred before the divergence of coelacanth and tetrapods (Fig. 3, branch b). Figs. 3 and 4 clearly show that amino acid replacements E122Q in RH1_{Lc} and in RH2_{Lc} occurred independently from each other. With the exception of RH2_{Lc} pigment, all RH1 and RH2 pigments have M207, whereas virtually all SWS1 and SWS2 pigments have L207. Thus, L207 is associated with the pigments with blue-shifted λ_{\max} values and M207L is suspected to cause a blue-shift in the λ_{\max} of RH2_{Lc} pigment. M207L occurred in RH2_{Lc} pigment after its divergence from the rest of RH2 pigments (Fig. 3, branch c). The inferred amino acids of the ancestral pigments support these amino acid replacements at branches b and c in Fig. 3. Mutants Q122E, L207M, and Q122E/L207M of RH2_{Lc} pigment have λ_{\max} values of 491 ± 1 nm, 484 ± 1 nm, and 499 ± 1 nm, respectively (Fig. 5). Thus, E122Q and M207L can cause a 21 nm blue-shift in the λ_{\max} and fully explain the adaptation of RH2_{Lc} pigment.

As already noted, the gecko RH2 pigment has a λ_{\max} value of 467 nm (Fig. 3). By constructing chimeric pigments, the cause of this blue-shift in the λ_{\max} was previously localized somewhere between transmembrane domains I and III (30).

The present analysis strongly suggests that E122Q in transmembrane domain III (Fig. 2) is a major cause of the blue shift in the λ_{\max} of the gecko RH2 pigment.

CONCLUSIONS

The molecular mechanisms of color vision in vertebrates may be elucidated in three steps: (i) cloning and characterization of the opsin genes; (ii) identification of amino acid changes that are potentially important in shifting the λ_{\max} of the pigments by using phylogenetic inferences; and (iii) determination of the actual effects of these mutations identified in the second step on the shifts in the λ_{\max} of visual pigments by using the *in vitro* assays (12–14). By so doing, we identified the potentially important amino acid replacements and tested specific hypotheses on the molecular mechanisms of adaptive evolution in the Comoran coelacanth.

The Comoran coelacanth evolved rod-rich retinas to adapt to a depth of about 200 m, where only a narrow range of colors at around 480 nm reaches from the surface. To visualize the entire range of available colors, coelacanth appears to have lost SWS1, SWS2, and LWS/MWS pigments and modified RH1_{Lc} and RH2_{Lc} pigments so that they can detect the two edges of the available light spectra, 478 nm and 485 nm. Our mutagenesis analyses based on this evolutionary hypothesis demonstrate that the coelacanth accomplished the coadaptations of RH1_{Lc} and RH2_{Lc} pigments to its unique habitat by two independent sets of amino acid replacements E122Q/A292S and E122Q/M207L, respectively.

We thank R. DeSalle for providing coelacanth tissue and J. Belote, P. Dunham, D. Frank, E. Maine, D. T. Sullivan, and R. Yokoyama for their comments on the manuscript. We also gratefully thank an anonymous reviewer for critically reviewing an earlier draft. This work was supported by National Institutes of Health Grant GM 42379.

1. Cloutier, R. & Ahlberg, P. E. (1996) in *Interrelationships of Fishes*, eds Stiassny, M. L., Parenti, L. R. & Johnson, G. D. (Academic Press, San Diego), pp. 445–479.
2. Maisey, J. G. (1996) *Discovering Fossil Fishes* (Henry Holt, New York).
3. Thompson, K. (1991) *Living Fossil: The Story of the Coelacanth* (Hutchinson, London).
4. Schliwien, U., Fricke, H., Scharl, M., Epplen, J. T. & Paabo, S. (1993) *Nature (London)* **363**, 405.
5. Fricke, H., Hissmann, K., Schauer, J. & Plante, R. (1995) *Nature (London)* **374**, 314.
6. Jerlov, N. G. (1976) *Marine Optics* (Elsevier, Amsterdam).
7. Locket, N. A. (1973) *Philos. Trans. R. Soc. London B* **266**, 493–521.
8. Locket, N. A. (1980) *Proc. R. Soc. London Ser. B* **208**, 265–307.
9. Dartnall, H. J. T. (1972) *Nature (London)* **239**, 341–342.
10. Wald, G. (1968) *Nature (London)* **219**, 800–807.
11. Yokoyama, S. (1994) *Mol. Biol. Evol.* **11**, 32–39.
12. Yokoyama, S. (1995) *Mol. Biol. Evol.* **12**, 53–61.
13. Yokoyama, S. (1997) *Annu. Rev. Genet.* **31**, 315–336.
14. Yokoyama, S. & Radlwimmer, F. B. (1998) *Mol. Biol. Evol.* **15**, 560–567.
15. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY), 2nd Ed.
16. Kawamura, S. & Yokoyama, S. (1998) *Vision Res.* **38**, 37–44.
17. Yokoyama, S., Radlwimmer, F. B. & Kawamura, S. (1998) *FEBS Lett.* **423**, 155–158.
18. Saitou, N. & Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406–425.
19. Felsenstein, J. (1985) *Evolution* **39**, 783–791.
20. Yang, Z. (1998) PAML, Phylogenetic Analysis by Maximum Likelihood (Univ. College London, London), Version 1.4.
21. Jones, D. T., Taylor, W. R. & Thornton, J. M. (1992) *Comput. Appl. Biosci.* **8**, 275–282.
22. Dayhoff, M. O., Schwartz, R. M. & Orcutt, B. C. (1978) in *Atlas of Protein Sequences and Structure*, ed. Dayhoff, M. O. (National Biomedical Research Foundation, Washington, DC), Vol. 5, pp. 345–352.

23. Nathans, J. & Hogness, D. S. (1983) *Cell* **34**, 807–814.
24. Nathans, J., Thomas, D. & Hogness, D. S. (1986) *Science* **232**, 193–202.
25. Zardoya, R. & Meyer, A. (1997) *Genetics* **146**, 995–1010.
26. Hope, A. J., Partridge, J. C., Dulai, K. S. & Hunt, D. M. (1997) *Proc. R. Soc. London Ser. B* **264**, 155–163.
27. Reitner, A., Sharpe, L. T. & Zrenner, E. (1991) *Nature (London)* **352**, 798–800.
28. Sakmar, T. P., Franke, R. R. & Khorana, H. G. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 8309–8313.
29. Sun, H., Macke, J. P. & Nathans, J. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 8860–8865.
30. Kojima, D., Oura, T., Hisatomi, O., Tokunaga, F., Fukada, Y., Yoshizawa, T. & Shichida, Y. (1996) *Biochemistry* **35**, 2625–2629.
31. Kawamura, S. & Yokoyama, S. (1995) *J. Mol. Evol.* **40**, 594–600.
32. Kawamura, S. & Yokoyama, S. (1996) *Vision Res.* **36**, 2797–2804.