

Skeleton to Hornworts & "Complex" Thalloid Liverworts

Revised through 17 May 2010

Group H – Hornworts			Group H
<i>Anthoceros</i> <i>Megaceros</i>	<i>Notothylas</i>	<i>Phaeoceros</i>	

Group C1 – Thalli rosette-like with embedded sporophytes			Group C1
<i>Oxymitra</i>	<i>Riccia</i> †	<i>Ricciocarpos</i>	

Group C2 – Thalli rolled into a blackish-purple tube when dry			Group C2
<i>Asterella</i> † <i>Mannia</i> †	<i>Plagiochasma</i> <i>Reboulia</i>	<i>Targionia</i>	

Group C3 – Thalli with gemmae receptacles			Group C3
<i>Lunularia</i>	<i>Marchantia</i>		

Group C4 – Thalli with distinctively protruding scales			Group C4
<i>Asterella</i> † <i>Athalamia</i>	<i>Mannia</i> † <i>Oxymitra</i>	<i>Riccia</i> †	

Group C5 – Thalli with dorsal air pores			Group C5
<i>Asterella</i> <i>Bucegia</i> <i>Conocephalum</i> <i>Corsinia</i> <i>Cryptomitrium</i>	<i>Lunularia</i> <i>Marchantia</i> <i>Mannia</i> <i>Peltolepis</i>	<i>Plagiochasma</i> <i>Preissia</i> <i>Reboulia</i> <i>Sauteria</i>	

Group C6 – Thalli lacking the above characteristics			Group C6
<i>Dumortiera</i>	<i>Riccia</i> †		

End.