

2007 No. 5

Large-Scale DNA Analysis of Bumble Bees: Worldwide Phylogeny Points to Commonality in the Decline of North American Species



Sydney Cameron, University of Illinois, Urbana-Champaign



A new phylogeny study sheds light on the relationship between different species of bumble bees throughout the world. This information helps narrow the search for which bumble bee species, important to agricultural crop pollination, are on the decline in the United States.









umble bees are among the most important wild bees in the United States. These insects are

increasingly used in the managed pollination of greenhouse crops, such as tomatoes and peppers. Despite their elevated importance to agriculture, bumble bee populations are on the decline in Western Europe and North America. Five bumble bee species in the United States are currently on the Xerces Society Red List of threatened insects. The timely work of entomology professor Sydney Cameron and her colleagues at the University of Illinois and the Natural History Museum in London contributes important insight into this critical agricultural problem.

BUMBLE BEE PHYLOGENY

Until recently, the relationship between bumble bee species remained unclear. Most prior bumble bee studies focused on morphological traits, which provided too few characters to resolve species relationships. In this study, the researchers conducted a phylogenetic analysis of nearly 90 percent of the world's 250 described bumble bee species based on DNA sequences from five genes, comprising approximately 4,000 base pairs. Independent Bayesian and parsimony analyses of each gene across 220 bumble bee species tested the reliability of the relationships estimated from the data. The researchers found strong evidence that the majority

Figure. 1. Images of bumble bees from the long-tongued group: a) *B.(Megabombus) supremus*; b) *B. (Diversobombus) trifasciatus*; and the short-tongued group c) *B. (Melanobombus) keriensis.*

Photograph Credit - Paul Williams, Natural History Museum, London.

* * * * *



Photograph Credit - James Whitfield, University of Illinois, Urbana-Champaign.

of bumble bee species fall into two distinct groups, the longertongued species in one group and their shorter-tongued sisters in the other. The complete bumble bee phylogeny is available as a PDF^{*}. Using the robust bumble bee phylogeny, Cameron's team inferred that bumble bees originated approximately 35 million years ago in central Asia and dispersed into the New World primarily between 15 and 10 million years ago.

Most bumble bees in the United States belong to the shorter-tongued group. From the phylogeny study, the researchers determined that out of the 47 bumble bee species in the United States, five of the declining species belong to the subgenus *Bombus sensu stricto*.

The reasons for the decline remain unclear and may vary regionally. Species in this subgenus include the most important pollinators for managed agriculture, specifically B. occidentalis in the United States and *B. terrestris* in Europe. *Bombus* occidentalis and another species B. franklini have all but disappeared in the United States. In addition, the range of their close relatives has been radically reduced compared to a decade ago. The bumble bee phylogenetic tree also reveals that two species, B. impatiens and B. vosnesenskii, within another subgenus, Pyrobombus, are undergoing range expansion in the United States. These species are distantly related to the species from the subgenus Bombus s.s., currently in decline.

Figure 1. Bumble bee field work on the Tibetan Plateau in the Sichuan basin of southwestern China.

IMPACT

The information gained from this novel and comprehensive phylogenetic tree may allow scientists to resolve the causes leading to the decline of subgenus *Bombus s.s*, and the increased range expansion in different bumble bees groups. Ultimately, the knowledge gained from understanding bumble bee relationships may assist scientists in working with other bumble bee groups to aid future agricultural applications.

*Bumble bee phylogeny available at - http://www.life.uiuc.edu/scameron/news/ USDAchronogram_maptree.pdf



The research reported in this fact sheet was sponsored by the Entomology and Nematology Program of the National Research Initiative Competitive Grants Program. To be placed on the mailing list for this publication or to receive additional information, please contact the NRI (202-401-5022 or NRICGP@ csrces.usda.gov). The fact sheet also is accessible via the NRI section of the Cooperative State Research, Education, and Extension Service Web site (http:// www.csrces.usda.gov/nri).

Any findings, conclusions, or opinions expressed by individuals in this research report are those of the authors and do not necessarily represent the policies of the U.S. Department of Agriculture. Publication of this fact sheet does not imply recommendation or endorsement by USDA over other research reports not mentioned. The U.S. Department of Agriculture (USDA), including CSREES, prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

For more information on civil rights and equal opportunity policies and programs, visit the <u>CSREES Equal Opportunity Office</u>.